


## Chemical Composition and Physiological Bioactivities of *Stiefia Multicaulis* (Vahl) Soják Vahl Grown in Palestine

Belal Rahhal<sup>1,\*</sup>Received: 1<sup>st</sup> May, 2024. Accepted 19<sup>th</sup> June, 2024, Published: 1<sup>st</sup> Feb, 2025, DOI: [10.35552/anjur.a.39.1.2266](https://doi.org/10.35552/anjur.a.39.1.2266)

**Abstract: Background:** Herbal products encompass a wide range of natural substances that show great potential as therapeutic materials. Consequently, there has been a substantial investigation into the therapeutic potential of numerous plant species in recent times. The present study sought to determine the components and assess the antioxidant, antilipase,  $\alpha$ -amylase, and  $\alpha$ -glucosidase enzyme inhibitory activities of *Stiefia multicaulis* (Vahl) Soják Vahl hexane, acetone, methanol, and aqueous fractions. **Methods:** Standard pharmacopeia methods were used to conduct screenings for both quantitative and qualitative phytoconstituents. The lipase,  $\alpha$ -amylase, and  $\alpha$ -glucosidase enzyme inhibitory activities were assessed using established reference methods. In addition, the antioxidant activity was assessed using a 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay. **Results:** The screening methods of pharmacopeias revealed that *S. multicaulis* contains a wide range of secondary metabolites, especially tannins. However, the most potent antioxidant, antilipase,  $\alpha$ -glucosidase, and  $\alpha$ -amylase inhibitory activities were acetone fraction, hexane, methanol, and aqueous *S. multicaulis* fractions, respectively. **Conclusion:** The current study outcomes are the first on the  $\alpha$ -amylase, lipase,  $\alpha$ -glucosidase inhibitory activities of *S. multicaulis* four solvent fractions. Further investigations are needed to assess the in vivo potential of these plant fractions in animal models.



**Keywords:** *Stiefia multicaulis*; Antioxidant; Antilipase;  $\alpha$ -Glucosidase;  $\alpha$ -Amylase.

### INTRODUCTION

As a result of its usefulness in the prevention and treatment of various diseases, phytomedicine has emerged as a staple of contemporary medicine. Their use as both conventional and complementary therapies dates back to ancient civilizations. Also, most modern medications are derived from plants because of the wide range of systems they can affect in the human body. Natural medicines and their derivatives are widely believed to be safe, but research has shown that they can have serious side effects or even be fatal if used improperly [1-4].

Recently, oxidative stress has become a widespread phenomenon. Once we have a lot of free radicals that exceed antioxidants, they can fight our body by many diseases, such as atherosclerosis, diabetes, cancer, Parkinson's, and Alzheimer's. Many scientists recommend avoiding the production of free radicals by minimizing their sources, such as stopping smoking and avoiding radiation and pollution [4, 5].

However, there is no way to avoid oxidative stress, but many antioxidants can help us, such as vitamin E and vitamin C supplements and beta-carotene supplements. In addition, a daily diet containing a variety of fruits and vegetables is also considered one of the methods used to obtain enough antioxidants [6].

Around 30% of the world's population is overweight, making obesity a global health epidemic, according to WHO biostatistical evaluations. On top of that, obesity is a major health problem in and of itself, and it often leads to other problems, including diabetes, heart disease, and gastrointestinal issues. The function of hyperlipidemia in the development of obesity has been the subject of recent research. In this regard, regulating obesity may

benefit from a reduction in dietary fat or its absorption through the inhibition of the lipase enzyme [7, 8].

People with diabetes mellitus (DM), a chronic, lifelong illness, experience hyperglycemia due to the body's incapacity to digest fat, carbs, and lipids. Over the last several decades, there has been a steady rise in the frequency of diabetes mellitus, with 422 million people worldwide living with the condition in 2014, up from 180 million in 1980. With 1.6 million deaths over the previous two years, it was the seventh most common cause of death, according to the World Health Organization [9-11].

*Stiefia multicaulis* (Vahl) Soják Vahl (family Lamiaceae) is an annual plant that reaches 30-50 cm. The leaves are long-petiolated, crenulate, ovate or oblong, obtuse, cordate or rounded at base, sometimes with two lateral minute segments; indumentum of mostly stellate dense appressed hairs which are denser still on the grayish lower face, while the floral leaves are sessile, membranous and oval. The calyx is 2-labiate, pale green or purple, glabrous with sessile glands, glandular-hairy or pilose. The origin is the west Irano-Turanian, extending into the Eastern regions of the Mediterranean basin. The major ingredients of *S. multicaulis* essential oil that are present in a high concentration are camphor, 1,8-cineole, borneol, and  $\alpha$ -pinene [12]. While it is often used in traditional medicine to treat colds, sore throats, and stomachaches, it has also been studied for its antispasmodic and antiseptic qualities [13].

The present study was undertaken to determine the components and assess the antioxidant, antilipase,  $\alpha$ -amylase, and  $\alpha$ -glucosidase enzyme inhibitory activities of *Stiefia multicaulis* (Vahl) Soják Vahl hexane, acetone, methanol, and aqueous fractions.

<sup>1</sup> Department of Biomedical Sciences, Faculty of Medicine and Health Sciences, An-Najah National University, Nablus, Palestine.

\* Correspondence: belalrahhal@najah.edu

## MATERIAL AND METHODS

**Instruments:** Spectrophotometer (Jenway 7315, England), Rotavap (Heidolph OB2000-VV2000, Germany), stir-mixer (Tuttnauer, Jerusalem), balance (Rad wag, AS 220/c/2 in Poland), freeze dryer (Mill rock technology-BT85, China), weighing scale (Adam Equipments, USA).

**Plant collection and preparation:** *S. multicaulis* leaves were collected in July 2018 from Wadi-Qana area in Nablus/Palestine. The plant was characterized in the Herbal Products Laboratory at An-Najah National University and kept within the voucher specimen code Pharm-PCT-2123.

The *S. multicaulis* leaves were rinsed with purified water multiple times before being left to dry in the shade at room temperature. The drying process took approximately one month, after which they were crushed roughly and stored in a tightly sealed glass jar until needed again.

**Fractionation method:** The dried powdered leaves were fractionated by four types of solvents: methanol, acetone, hexane, and distilled water. The dried leaves were soaked in 1 liter of each solvent sequentially and each fraction was placed in a shaker device for about two days at room temperature. The filtrates were concentrated under a vacuum on a rotary evaporator or dried using a freeze drier. The final fractions were stored in a refrigerator [14].

**Phytochemical analysis:** Each fraction (methanol, acetone, hexane, and water) underwent qualitative chemical analysis for each primary and secondary metabolic compound such as steroids, terpenoids, starch, carbohydrates, cardiac glycosides, flavonoids, tannins, monosaccharide, protein, reducing sugar, alkaloids, phenols, and saponins. Phytochemical analysis was executed according to the standard qualitative analytical assays as described by Harborne and Evans [15, 16].

**Quantitative total phenol content:** Total phenol content was determined in the four plant fractions using the spectrophotometric and Folin-Ciocalteu's reagent. In milligrams of gallic acid, equivalent per gram of plant fractions is the concentration that was determined from the absorbance measurements [17].

**Determination of total tannin content:** Its total amount of tannin was ascertained using a modified Folin-Denis colorimetric technique. The calibration curve will be constructed by making serial dilutions from an aqueous stock solution of 1 mg/ml tannic acid. Catechin equivalent concentrations (mg CAE/g) for all plant components.

**Porcine pancreatic lipase enzyme inhibitory method:** By dissolving 100 mg of each plant fraction in 100 ml of 10% DMSO, a working solution (1 mg/ml) was prepared for each *S. multicaulis* fraction. Subsequently, the solution was diluted to reach various concentrations (0.05, 0.1, 0.2, 0.3, and 0.4 mg/ml). Be that how it may, the following equation was used to determine the lipase enzyme inhibitory potential (%) [18]:

$$I (\%) = \frac{[ABS_{blank} - ABS_{test}]}{[ABS_{blank}]} * 100\%$$

**$\alpha$ -Amylase inhibitory method:** To make a working solution (1 mg/ml) from each *S. multicaulis* plant fraction, 25 mg of each fraction was dissolved in a small quantity of 10% DMSO. To this, 25 ml of a buffer solution was added. Various dilutions (0.01, 0.05, 0.07, 0.1, and 0.5 mg/ml) were obtained using the buffer. The absorbance was measured at 540 nm using a UV-Vis Spectrophotometer, with Acarbose serving as a positive

reference solution. The inhibition ability against  $\alpha$ -amylase was determined using a specific equation.

$$I (\%) = \frac{[ABS_{blank} - ABS_{test}]}{[ABS_{blank}]} * 100\%$$

Where I (%), is the  $\alpha$ -amylase inhibitory percentage [19].

**$\alpha$ -Glucosidase inhibitory activity:** One milligram per milliliter of phosphate buffer was used to create a workable solution from 100 milligrams of each *S. multicaulis* leaf portion. Phosphate buffer diluted the solution to get different concentrations (100, 200, 300, 400, 500  $\mu$ g/ml). The next calculation was used to assess the inhibiting action of  $\alpha$ -glucosidase.

$$I (\%) = \frac{[ABS_{blank} - ABS_{test}]}{[ABS_{blank}]} * 100\%$$

Where I (%), is the percentage inhibition of  $\alpha$ -glucosidase [20].

**Free radical scavenging activity:** A mixture containing 1 mg/ml of each *S. multicaulis* leaf fragment was made by dissolving 100 mg of each fraction in 100 ml of methanol. Subsequently, the solution was further diluted with methanol to achieve various concentrations ranging from 2 to 100  $\mu$ g/ml. A UV-Vis spectrophotometer measured the absorbance at 517 nm, with Trolox functioning as a control. The results were contrasted with the control. The subsequent formula was used to compute the antioxidant activity.

$$I (\%) = \frac{[ABS_{blank} - ABS_{test}]}{[ABS_{blank}]} * 100\%$$

where I (%), is the percentage of antioxidant activity [21, 22].

**Statistical analysis:** All tests were repeated three times, and the obtained results are shown as means  $\pm$  SD.

## RESULTS AND DISCUSSION

**Phytochemical analysis:** Table 1 shows the phytochemical identification test outcomes, which revealed that *S. multicaulis* all fractions containing tannins, while the acetone, hexane and methanol fractions contain glycosides. Moreover, the aqueous fraction was the only one containing phenol.

**Table (1):** The chemical analysis of *S. multicaulis* extracts.

Plant products	Acetone	Hexane	Methanol	Water
Protein	-	-	+	+
Carbohydrate	-	-	+	+
Tannin	+	+	+	+
Flavonoid	-	-	-	-
Saponin	-	-	-	-
Glycosides	+	+	+	-
Phenol	-	-	-	+


**Total tannin content:** Estimations of total tannin contents in *S. multicaulis* leave methanol, hexane, acetone and aqueous fractions were conducted according to standard analytical method. As well as catechin was used as a reference compound while the absorbance was measured at  $\lambda_{max}$  =500 nm. However, from the standard calibration curve (Fig. 1), the total tannin content in all plant fractions was calculated according to the following formula:

$$Y = 0.001X + 0.002, R^2 = 0.991, \text{ Where}$$

Y- Absorbance at 500 nm

X- Total tannin in the plant fraction.

The quantitative tannin contents of *S. multicaulis* leave methanol, hexane, acetone, and aqueous fractions are presented in Table 2 and showed that the aqueous fraction has the highest percent of tannin contents compared with other fractions.



**Figure (1):** Standard calibration curve of Catechin.


**Total phenol contents:** Estimating total phenols in *S. multicaulis* leaves was conducted in the aqueous fraction, while the other fractions did not contain phenolic compounds. The total phenols were expressed as mg/g Gallic acid equivalent and the standard calibration curve of Gallic acid (Fig. 2) was constructed

to estimate the total phenol content in the plant aqueous fraction using the following formula:

$$Y = 0.009X + 0.021, R^2 = 0.992, \text{ Where}$$

Y- Absorbance at 760 nm

X- Total phenol in the aqueous fraction.



**Figure (2):** Standard calibration curve of Gallic acid

However, quantitative phenol contents of *S. multicaulis* leave aqueous fraction presented in Table 2 and demonstrated that this fraction contains 14.64±1.53 GAE/g of dry fraction.

**Table (2):** Quantitative total phenols, and tannins contents of *S. multicaulis* leave hexane, acetone, methanol, and aqueous fractions.


Plant fractions	Total Tannin contents, mg of CAE/g of dry fraction, ±SD	Total phenol contents, mg of GAE/g of dry fraction, ±SD
Hexane	3.95±0.59	-
Acetone	1.54±0.61	-
Methanol	3.95±0.59	-
Aqueous	9.25±2.25	14.64±1.53

**Antioxidant activity:** As observed in Table 3, the results revealed that *S. multicaulis* four solvents fractions have free radical scavenging property in dose-dependent manner (Fig. 3)

and all the screened fractions have potent antioxidant activity as presented in Table 3.

**Table (3):** The IC<sub>50</sub> values and DPPH inhibitory activity by *S. multicaulis* four solvents fractions and Trolox.

Conc.	Trolox ±SD	Hexane fraction ±SD	Acetone fraction ±SD	Methanol fraction ±SD	Aqueous fraction ±SD
0	0±0	0±0	0±0	0±0	0±0
2	53.89±3.6	55.47±0.53	59.99±0.53	36.98±0.53	23.21±0.44
5	94.15±0.80	55.47±0.53	61.13±0.53	42.07±0.8	39.84±1.86
10	96.75±1.80	57.35±0.53	65.84±0.26	56.98±1.06	49.4±0.41
20	97.4±1.80	61.5±0.53	73.2±0.01	83.39±0.53	49.4±0.41
50	97.72±2.20	96.79±0.26	76.98±0.53	92.26±0.26	91.69±1.06
100	99.35±0.00	97.73±0.34	82.45±0.80	93.95±0.53	92.45±0.29
IC <sub>50</sub> (µg/ml) ±SD	2.23±1.57	5.37±0.40	4.78±0.44	6.60±0.62	10.23±0.7




**Figure (3):** DPPH scavenging properties by *S. multicaulis* four solvents fractions and Trolox.

The antioxidant outcomes revealed that *S. multicaulis* hexane, methanol, acetone and aqueous solvents fractions have potential antioxidant activity while the acetone fraction has the highest antioxidant potential with IC<sub>50</sub> value of 4.78±0.44 µg/ml in comparison with Trolox a potential free radical scavenging compound and vitamin E analogue which has antioxidant IC<sub>50</sub> value of 2.23±1.57 µg/ml.

**α-Amylase inhibitory activity:** Figure 4 depicted the α-amylase inhibitory activity of *S. multicaulis* four fractions in comparison with Acarbose, which is used therapeutically in the management of type 3 of diabetes. Moreover, Table 4 shows the inhibition percentages and the IC<sub>50</sub> values of all the studied samples. However, the aqueous fraction showed the highest α-amylase inhibitory activity with an IC<sub>50</sub> value of 158.48±0.62 µg/ml.

**Table (4):** The α-amylase inhibitory activity by Acarbose drug and *S. multicaulis* four solvents fractions in addition to their IC<sub>50</sub> values.

Conc.	Acarbose ±SD	Hexane fraction ±SD	Acetone fraction ±SD	Methanol fraction ±SD	Aqueous fraction ±SD
0	0±0	0±0	0±0	0±0	0±0
10	53.22±1.20	8.3±0.24	3.89±0.72	18.47±0.24	38.13±1.20
50	54.91±0.58	10.84±0.96	23.04±0.95	23.38±0.47	38.81±0.24
70	66.1±1.34	24.74±0.96	25.25±0.24	37.28±1.43	61.35±0.96
100	66.1±1.62	24.74±0.96	29.82±0.95	43.04±0.47	63.55±0.24
500	72.54±1.37	26.43±0.95	60.67±1.43	51.69±1.20	65.76±0.48
IC <sub>50</sub> (µg/ml) ±SD	28.84±1.22	50118±0.81	501.2±0.86	398.1±0.76	158.48±0.62




**Figure (4):**  $\alpha$ -Amylase inhibitory activity of *S. multicaulis* four solvents fractions and Acarbose drug.

**$\alpha$ -Glucosidase inhibitory activity:** Alongside the reference medication Acarbose,  $\alpha$ -glucosidase inhibitory activity was investigated for *S. multicaulis* four solvent fractions. When compared to the positive control, Acarbose, which has an  $IC_{50}$

value of  $37.15 \pm 0.33$   $\mu\text{g/ml}$ , the findings indicated that the methanolic fraction had the strongest  $\alpha$ -glucosidase inhibitory activity (Fig. 5 and Table 5).

**Table (5)** The  $\alpha$ -glucosidase suppressant action by Acarbose drug and *S. multicaulis* four solvents fractions in addition to their  $IC_{50}$  doses.

Conc.	Acarbose $\pm$ SD	Hexane fraction $\pm$ SD	Acetone fraction $\pm$ SD	Methanol fraction $\pm$ SD	Aqueous fraction $\pm$ SD
0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0
100	65.8 $\pm$ 0.42	17.49 $\pm$ 0.20	25.73 $\pm$ 1.03	65.29 $\pm$ 0.41	23.21 $\pm$ 0.44
200	67.75 $\pm$ 0.35	20.29 $\pm$ 0.41	32.2 $\pm$ 0.21	66.02 $\pm$ 1.44	39.84 $\pm$ 1.86
300	73.20 $\pm$ 0.42	22.2 $\pm$ 0.62	35.87 $\pm$ 0.41	70.43 $\pm$ 0.20	49.4 $\pm$ 0.41
400	85.35 $\pm$ 0.35	37.64 $\pm$ 0.41	66.02 $\pm$ 1.03	72.35 $\pm$ 0.83	49.4 $\pm$ 0.41
500	92.22 $\pm$ 0.11	38.96 $\pm$ 0.62	66.02 $\pm$ 1.03	89.11 $\pm$ 0.82	65.44 $\pm$ 0.62
$IC_{50}$ ( $\mu\text{g/ml}$ ) $\pm$ SD	37.15 $\pm$ 0.33	12589 $\pm$ 0.45	389 $\pm$ 0.74	39.81 $\pm$ 0.74	316.22 $\pm$ 0.75




**Figure (5):**  $\alpha$ -Glucosidase inhibitory potentials by Acarbose drug and *S. multicaulis* four solvents fractions.

**Antilipase activity:** We used the pig pancreatic lipase inhibitory test to assess the antilipase activity of the four fractions of the *S. multicaulis* plant. We compared the results to those of the commercial medication Orlistat, which is a positive control.

Table 6 and Fig. 6 demonstrated that, compared to the positive control Orlistat, which had an  $IC_{50}$  value of  $12.3 \pm 0.33 \mu\text{g/ml}$ , the hexane fraction exhibited the strongest antilipase activity at  $316.2 \pm 0.87 \mu\text{g/ml}$ .

**Table (6):** The lipase inhibitory activity by Orlistat drug and *S. multicaulis* four solvents fractions in addition to their  $IC_{50}$  values.

Conc.	Orlistat $\pm$ SD	Hexane fraction $\pm$ SD	Acetone fraction $\pm$ SD	Methanol fraction $\pm$ SD	Aqueous fraction $\pm$ SD
0	$0 \pm 0$	$0 \pm 0$	$0 \pm 0$	$0 \pm 0$	$0 \pm 0$
50	$51.65 \pm 0.21$	$22.49 \pm 1.52$	$11.54 \pm 0.5$	$26.9 \pm 0.67$	$13.68 \pm 0.5$
100	$64.80 \pm 0.42$	$30.59 \pm 0.5$	$11.54 \pm 0.5$	$26.9 \pm 0.67$	$13.68 \pm 0.5$
200	$78.13 \pm 0.19$	$35.59 \pm 0.16$	$20.23 \pm 8.41$	$42.25 \pm 0.84$	$20.94 \pm 0.67$
300	$85.72 \pm 0.53$	$51.3 \pm 1.52$	$40.23 \pm 0.16$	$42.25 \pm 0.84$	$32.49 \pm 0.5$
400	$94.67 \pm 0.31$	$72.13 \pm 0.67$	$42.73 \pm 0.16$	$52.14 \pm 0.33$	$48.8 \pm 1.68$
$IC_{50} (\mu\text{g/ml}) \pm$ SD	$12.3 \pm 0.33$	$316.2 \pm 0.87$	$6309 \pm 1.95$	$630.9 \pm 0.67$	$645.6 \pm 0.77$



**Figure (6):** Lipase inhibitory property by Orlistat drug and *S. multicaulis* four solvents fractions.

To the best of the author's knowledge, no previous investigations evaluated the  $\alpha$ -amylase, lipase, and  $\alpha$ -glycosidase inhibitory activities and the current study will be the first one conducted these experiments.

## DISCUSSION

Medicinal plants and phytochemicals have been used in traditional medicine worldwide for the treatment of many ailments since ancient times. Crude plant extracts are now recognized as a significant reservoir of natural compounds that may be used in the creation of medications to combat a range of ailments, the formulation of pharmaceutical goods, and innovative biomedical research.

*Stefia multicaulis* (synonym *Salvia multicaulis* Vahl) is one of the traditional medicinal plants in Palestine and other Mediterranean regions [24].

Antioxidants are essential in defending against oxidative stress and the harm caused by free radicals, which are triggered by diabetes, cardiovascular diseases, and cancer. The DPPH free radicals screening showed that all *S. multicaulis* hexane, acetone, methanol, and aqueous fractions have potent

antioxidant effects compared with Trolox with  $IC_{50}$  doses of  $5.37 \pm 0.40$ ,  $4.78 \pm 0.44$ ,  $6.60 \pm 0.62$ ,  $10.23 \pm 0.7$ , and  $2.23 \pm 1.57 \mu\text{g/ml}$ , respectively. These outcomes agree with a study conducted by Tepe *et al.*, which showed that the essential oil and the methanolic extract of *S. multicaulis* have antioxidant activities with  $IC_{50}$  values of  $2.4 \pm 0.05$  and  $16.3 \pm 0.1 \mu\text{g/ml}$ , respectively [23]. Besides, these results agree with Pehlivan and Sevindik investigation, which found that the *S. multicaulis* ethanolic extract had a potent antioxidant effect [25].

However, our phytochemical screenings showed that the *S. multicaulis* enriches phytochemicals; all extracts also have tannins, and the aqueous extract contains a high number of phenols. These components, including tannins and phenols, have high potential antioxidant effects [26].

Moreover, the aqueous *S. multicaulis* fraction showed the highest  $\alpha$ -amylase inhibitory activity with an  $IC_{50}$  value of  $158.48 \pm 0.62 \mu\text{g/ml}$ . This outcome may be due to the richness of *S. multicaulis* aqueous fraction with tannins and phenols ( $9.25 \pm 2.25 \text{ mg}$  of CAE/g of dry fraction and  $14.64 \pm 1.53 \text{ mg}$  of GAE/g of dry fraction, respectively).

In fact, tannins and phenols are among the most potent anti-amylase agents [27]. Compared to the positive control, Acarbose, which has an IC<sub>50</sub> value of 37.15±0.33 µg/ml, the findings indicated that the methanolic fraction had the strongest α-glucosidase inhibitory activity (IC<sub>50</sub>=39.81±0.74 µg/ml). These findings agree with the study by Tamimi et al., which found that plants rich in phytochemicals had potential antiglycosidase effects [28].

*S. multicaulis* hexane, acetone, methanol, and aqueous fractions revealed mild lipase inhibitory properties compared with Orlistat drug and the hexane fraction showed the highest antilipase effect with IC<sub>50</sub> dose of 316.2±0.87 µg/ml while Orlistat has antilipase effect IC<sub>50</sub> dose of 12.3 ±0.33 µg/ml.

Tannins and phenolic molecules, which are found in many crops, have substantial inhibitory effects on digestive enzymes such as lipase, α-glucosidase, and α-amylase. The bioactive substances can attach to these enzymes, which leads to a decrease in their activity and, therefore, reduces the absorption of fats and carbs. This inhibitory process is essential for controlling metabolic diseases such as obesity and diabetes by reducing fat absorption and regulating blood glucose levels after a meal. The mentioned components are derived from green tea, grapes, pomegranates, and nuts and seeds. Their ability to act as both enzyme inhibitors and antioxidants further amplifies their potential health advantages. Therefore, including tannins and phenols in one's diet may provide a natural approach to enhancing metabolic health and controlling chronic illnesses [29].

In fact, to the best of our knowledge, no previous investigations screened the antilipase, antiglycosidase and anti-amylase effects of *S. multicaulis* four fractions.

#### Limitations

The obtained data were done in vitro from crude extract, future studies will undertake to study the effects of this plant in vivo from pure compounds.

#### CONCLUSION

The obtained data about the *S. multicaulis* aqueous, hexane, methanol, and acetone solvent fractions showed that they have potential antioxidant potentials compared with Trolox. In addition, it can be considered as the first information on the α-amylase, lipase and α-glycosidase inhibitory activities of *S. multicaulis* four solvent fractions. Further investigations are needed to assess the in vivo potential of these plant fractions in animal models.

#### DISCLOSURE STATEMENT

- Ethics approval and consent to participate; Not applicable.
- Consent for publication: Not applicable
- Availability of data and materials: This published article and its supplementary information files include all data generated or analyzed during this study.
- Author's contribution: BR: Conceptualization, Validation, Investigation, writing – original draft, Writing - review & editing, Visualization, analysis, Supervision, Project administration.
- Funding: None
- Conflicts of interest: The authors declare that they have no competing interests.
- Acknowledgements: The author is thankful for Mss Linda Issa for her technical assistance.

#### REFERENCES

- 1] Ukrainets I, Sidorenko L, Gorokhova O, Jaradat N. 4-Hydroxy-2-quinolones. 93. Synthesis and biological properties of 2-hydroxy-4-imino-1, 4-dihydroquinoline-3-carboxylic acid NR-amides. *Chemistry of Heterocyclic Compounds*. 2006;42:475-87.
- 2] Shawahna R, Jaradat NA. Ethnopharmacological survey of medicinal plants used by patients with psoriasis in the West

Bank of Palestine. *BMC complementary and alternative medicine*. 2017;17:1-9.

- 3] Jaradat NA. Review of the taxonomy, ethnobotany, phytochemistry, phytotherapy and phytotoxicity of germander plant (*Teucrium polium* L.). *medicine*. 2015;3(4).
- 4] Qneibi M, Jaradat N, Hawash M, Olgac A, Emwas N. Ortho versus meta chlorophenyl-2, 3-benzodiazepine analogues: synthesis, molecular modeling, and biological activity as AMPAR antagonists. *ACS omega*. 2020;5(7):3588-95.
- 5] Saeidnia S, Abdollahi M. Toxicological and pharmacological concerns on oxidative stress and related diseases. *Toxicology and applied pharmacology*. 2013;273(3):442-55.
- 6] Poljsak B. Strategies for reducing or preventing the generation of oxidative stress. *Oxidative medicine and cellular longevity*. 2011;2011.
- 7] Hurt RT, Kulisek C, Buchanan LA, McClave SA. The obesity epidemic: challenges, health initiatives, and implications for gastroenterologists. *Gastroenterology & hepatology*. 2010;6(12):780.
- 8] James PT. Obesity: the worldwide epidemic. *Clinics in dermatology*. 2004;22(4):276-80.
- 9] Collaboration ERF. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *The Lancet*. 2010;375(9733):2215-22.
- 10] Mrabti HN, Sayah K, Jaradat N, Kichou F, Ed-Dra A, Belarj B, et al. Antidiabetic and protective effects of the aqueous extract of *Arbutus unedo* L. in streptozotocin-nicotinamide-induced diabetic mice. *Journal of Complementary and Integrative Medicine*. 2018;15(3).
- 11] NCD Risk Factor Collaboration. Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4. 4 million participants. *The Lancet*. 2016;387(10027):1513-30.
- 12] Morteza-Semnani K, Moshiri K, Akbarzadeh M. The essential oil composition of *Salvia multicaulis* Vahl. *Journal of Essential Oil Bearing Plants*. 2005;8(1):6-10.
- 13] Erdemoglu N, Turan NN, Caköcö I, Sener B, Aydön A. Antioxidant activities of some Lamiaceae plant extracts. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*. 2006;20(1):9-13.
- 14] Jaradat N, Al-Masri M, Al-Rimawi F, Naser A, Zaid MMS, Hussein F, et al. Investigating the impacts of various solvent fractions of *Bupleurum lancifolium* on the anti-microbial and antioxidant potentials. *JOURNAL OF INTERCULTURAL ETHNOPHARMACOLOGY*. 2017;6(4):401-6.
- 15] Evans WC. *Trease and Evans Pharmacognosy, International Edition E-Book*. USA: Elsevier Health Sciences; 2009.
- 16] Harborne A. *Phytochemical methods a guide to modern techniques of plant analysis*. USA: Springer Science & Business Media; 1998.
- 17] Jaradat N, Hussen F, Al Ali A, Alniss H, Dweikat M. Phytoconstituents, free radical scavenging potential, total phenols and total flavonoids assessments for Violet Horned Poppy from Jerusalem Mountains. *J Mater Environ Sci*. 2015;6(10):2958-66.
- 18] Jaradat N, Zaid A, Hussein F, Zaqqouq M, Aljammal H, Ayesh O. Anti-lipase potential of the organic and aqueous extracts of ten traditional edible and medicinal plants in Palestine; a comparison study with orlistat. *Medicines*. 2017;4(4):89.
- 19] Ali H, Houghton P, Soumyanath A. α-Amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*. *Journal of ethnopharmacology*. 2006;107(3):449-55.

- 20] Kim J-S, Kwon C-S, SoN KH. Inhibition of alpha-glucosidase and amylase by luteolin, a flavonoid. *Bioscience, biotechnology, and biochemistry*. 2000;64(11):2458-61.
- 21] Jaradat NA, Abualhasan M. Comparison of phytoconstituents, total phenol contents and free radical scavenging capacities between four Arum species from Jerusalem and Bethlehem. *Pharmaceutical Sciences*. 2016;22(2):120.
- 22] Hawash M, Eid AM, Jaradat N, Abualhasan M, Amer J, Naser Zaid A, et al. Synthesis and biological evaluation of benzodioxole derivatives as potential anticancer and antioxidant agents. *Heterocyclic Communications*. 2020;26(1):157-67.
- 23] Tepe B, Donmez E, Unlu M, Candan F, Daferera D, Vardar-Unlu G, et al. Antimicrobial and antioxidative activities of the essential oils and methanol extracts of *Salvia cryptantha* (Montbret et Aucher ex Benth.) and *Salvia multicaulis* (Vahl). *Food chemistry*. 2004;84(4):519-25.
- 24] Barhoumi LM, Al-Jaber HI, Abu Zarga MH. A new diterpene and other constituents of *Salvia multicaulis* from Jordan. *Natural Product Research*. 2022;36(19):4921-8.
- 25] Pehlivan M, Sevindik M. Antioxidant and antimicrobial activities of *Salvia multicaulis*. *Turkish Journal of Agriculture-Food Science and Technology*. 2018;6(5):628-31.
- 26] Jaradat NA, Shawahna R, Hussein F, Al-Lahham S. Analysis of the antioxidant potential in aerial parts of *Trigonella arabica* and *Trigonella berythea* grown widely in Palestine: a comparative study. *European Journal of Integrative Medicine*. 2016;8(5):623-30.
- 27] Jaradat N, Dwikat M, Amer J, Ghanim M, Hawash M, Hussein F, et al. Total phenolic contents, cytotoxic, free radicals, porcine pancreatic  $\alpha$ -amylase, and lipase suppressant activities of *Artemisia dracunculoides* plant from Palestine. *Frontiers in Pharmacology*. 2024;15:1351743.
- 28] Rabi L, Altamimi M, Jaradat N. Phytoconstituents, antioxidant and inhibitory activity against  $\alpha$ -amylase and  $\alpha$ -glucosidase of *Opuntia ficus-indica*. *An-Najah University Journal for Research-A (Natural Sciences)*. 2020;34(1):31-48.
- 29] Jaradat NA, Zaid AN, Abuzant A, Khalaf S, Abu-Hassan N. Phytochemical and biological properties of four *Astragalus* species commonly used in traditional Palestinian medicine. *European Journal of Integrative Medicine*. 2017;9:1-8.