

## Validating An Instrument for Measuring the Awareness and Challenges of Nursing Services Accreditation from the Healthcare Providers in Yemen

Talal Mansoor<sup>1,2,\*</sup>, Malakeh.Z. Malak<sup>2</sup>, Sharifa Ezat Bint Wan Puteh<sup>3</sup>, Azimatun Noor Aizuddin<sup>1</sup> & Nawal Saeed Banafa<sup>2</sup>

(Type: Full Article). Received: 28<sup>th</sup> Sep. 2024. Accepted: 27<sup>th</sup> Nov. 2024, Published: 1<sup>st</sup> Oct. 2025,  
DOI: <https://doi.org/10.59049/2790-0231.10.4.2391>

(This article belongs to the Special Issue: Trends in Nursing)

**ABSTRACT: Introduction:** Enhancing healthcare quality and patient safety is a global priority. Unsafe healthcare practices increase patient morbidity, mortality, and suffering while undermining public trust in healthcare systems. This study aims to validate an instrument assessing awareness and challenges related to nursing services accreditation among healthcare providers (HCPs) in Yemen, supporting broader efforts to improve healthcare quality and patient safety. **Materials and Methods:** Using stratified random sampling, 358 healthcare providers (nurses and doctors) were selected from various hospitals in Hadhramaut, Yemen. Structural Equation Modeling with Confirmatory Factor Analysis (CFA) was employed to validate the measurement model. Key metrics for validity and reliability, including convergent validity (average variance extracted [AVE] and composite reliability [CR]), discriminant validity, internal consistency (Cronbach's Alpha), and goodness-of-fit (GOF) indices, were rigorously assessed to confirm model robustness. **Results:** The validated instrument included nine constructs and 39 items. The CFA demonstrated excellent model fit, with all factor loadings exceeding 0.50, affirming strong convergent validity. Discriminant validity was achieved as each factor's AVE exceeded the squared correlations with other factors. Reliability measures were met, with Cronbach's Alpha values above 0.70, CR above 0.60, and AVE above 0.50 for all constructs. **Conclusion:** This study successfully validated an instrument for assessing nursing services accreditation awareness among healthcare providers in Hadhramaut hospitals, Yemen. This reliable and valid tool provides a framework for evaluating accreditation awareness and can be adapted to similar healthcare settings. The findings offer practical insights for healthcare professionals, researchers, and policymakers to guide quality and safety improvements in healthcare.

**Keywords:** Healthcare Providers, Awareness, Measurement Model, Confirmatory Factor Analysis, Hadhramaut, Yemen.

### INTRODUCTION

Accreditation is a widely recognized mechanism for managing and evaluating healthcare quality globally. It not only enhances overall healthcare services but also strengthens nursing practice, promoting autonomy in patient care, professional communication within multidisciplinary teams, collaboration, professional development, and decentralized decision-making [1,2]. Nursing plays a crucial role in healthcare systems, directly influencing healthcare outcomes [3]. Nurses significantly contribute to patient outcomes and care quality, ensuring patient safety [1, 4].

However, the successful adoption of accreditation in healthcare settings is often hindered by a lack of awareness and understanding among healthcare professionals, as well as insufficient commitment from management and physicians [5,6]. Limited knowledge about accreditation processes can severely impact effectiveness, leading to suboptimal outcomes [7].

Furthermore, studies have reported inconsistent findings regarding the impact of accreditation on healthcare organizational performance and patient satisfaction [8]. It is vital for hospital staff to grasp the comprehensive nature and importance of accreditation, as this understanding can lead to

improved care quality when implemented correctly [9].

In Yemen, the healthcare system faces significant challenges, including low staff morale, inadequate service quality, and a fragile healthcare infrastructure [10-12]. This is particularly evident in Hadramaut, characterized by diverse geography, including coastal areas, mountains, and valleys [13]. With a population of approximately 1.5 million, Hadramaut has a mix of urban and rural communities, affecting access to healthcare services. Urban centers like Mukalla have better facilities, while rural areas often suffer from a shortage of essential services, leading to disparities in health outcomes [13,14].

Healthcare services in Hadramaut are strained by various factors, including a lack of trained professionals, inadequate facilities, and limited access to essential medications [15,16]. The region has also been adversely affected by ongoing political instability and economic challenges, exacerbating health issues such as maternal and child morbidity and communicable and non-communicable diseases [11,14,16]. The absence of standardized accreditation and adherence to international best practices has hindered Yemen's progress in healthcare standards compared to other developing countries.

1 Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur.

2 Department of Community Health, Faculty of Nursing, Hadhramout University, Hadhramout, Yemen.

\*Corresponding author email: [raed@zu.edu.jo](mailto:raed@zu.edu.jo)

3 Community Health Nursing, Faculty of Nursing, Al-Zaytoonah University of Jordan, Amman, Jordan.

Despite global efforts to enhance healthcare quality through accreditation, a significant gap remains in understanding how healthcare providers, especially in resource-limited and politically unstable regions like Yemen, perceive and engage with accreditation processes [17,18]. Addressing this gap is crucial, as accreditation can catalyze improvements in healthcare quality, patient safety, and overall system performance, even in challenging environments. Moreover, there is limited understanding of the awareness of nursing service accreditation standards among healthcare providers, including decision-makers, in Yemen [8,18,19].

### Purpose of study

The aim of this study is to validate an instrument designed to assess awareness and challenges related to nursing services accreditation among healthcare providers in Yemen, contributing to broader efforts to enhance healthcare quality and patient safety.

## MATERIALS AND METHODS

### Design

A descriptive cross-sectional correlational design was employed.

### Sample and setting

The research focused on healthcare professionals (HCPs) working in hospitals in Hadramaut, including physicians, nurses, and administrators. A total of 394 individuals were sampled from eight hospitals—four public hospitals (two governorates and two district) and four private hospitals (two for-profit and two non-profit). The selection ensured representation from both public and private sectors.

To select hospitals, we employed simple probability sampling from a comprehensive list of eligible hospitals in Hadramaut. Hospitals were categorized based on type and operational characteristics, with two hospitals chosen from each category to reflect the diversity of healthcare services available in the region.

Using OpenEpi software, the initial sample size for healthcare professionals was calculated at 358 based on the study's descriptive nature (Table 1). To mitigate potential information collection bias, the sample size was increased by 10%, resulting in a final sample size of 394 participants. This sample was allocated to selected hospitals according to the percentage of healthcare professionals present in each category (nurses, doctors, and administrators). Random sampling within each hospital was conducted, allowing participants to respond to surveys at their convenience.

**Table (1):** Distribution of Sample Size among the Selected Hospitals.

No.	Hospital	Percentage	Sample size distribution
1	Ibn Sina Teaching hospital	23%	82
2	Seyun General hospital	22%	79
3	Alsheher hospital	19%	68
4	Al-Qatn General Hospital	10%	36
5	Al Bourj hospital Advisory	9%	32
6	Bin Zeela Modern Hospital	8%	29
7	Dawan Hospital-khailatBugshan	6%	21
8	Saleh Babeker Charity Hospital	3%	11
	Total	100%	358

### Instruments

The quantitative data collection instrument for this study was a self-reporting questionnaire, which underwent pre-testing to

ensure reliability and validity. Approval for the questionnaire was granted by two lecturers at Hadramaut University. To establish its validity and reliability, a review by an expert and conducting a pilot study were implemented. The instrument underwent review by two experts, and a pilot test was conducted with 36 healthcare providers (HCPs), including nurses, doctors, and managers, as a pre-test sample. Subsequently, the instrument's reliability was assessed and validated using SPSS software.

Once approved, the questionnaire was distributed to the available HCPs. Detailed instructions for completing the questionnaire were provided on the first page. The questionnaires were personally given to the respondents at the beginning of their morning shift and collected at the end of their shift. This process was continuous until the essential sample for each hospital was obtained without any data duplication.

The questionnaire comprised three parts. **Part 1:** HCPs' personal sociodemographic, occupational, and organizational data, designed based on the literature review.

**Part 2:** The Awareness Regarding Accreditation (ARA) tool is a structured questionnaire comprising 13 items designed to assess healthcare professionals' awareness and understanding of accreditation processes and standards. It evaluates several domains, including general awareness, knowledge of accreditation standards, perceived benefits, challenges to implementation, and attitudes toward accreditation. Each item typically uses a Likert scale for responses, ranging from "Strongly Agree" to "Strongly Disagree," allowing for a nuanced understanding of participants' perceptions. The scoring system involves assigning numerical values to responses (e.g., 1 for "Strongly Disagree" to 5 for "Strongly Agree"), with higher total scores indicating greater awareness and understanding of accreditation. This quantitative data can be analyzed to identify knowledge gaps and inform targeted educational interventions to enhance accreditation engagement among healthcare professionals.

**Part 3:** The Awareness of Nursing Services Accreditation Standards tool is a comprehensive assessment instrument that evaluates healthcare professionals' knowledge across five key domains derived from the Malaysian Society for Quality in Health (MSQH) standards: Organization and Management (OM), Human Resource Development and Management (HRD), Policies and Procedures (PP), Facilities and Equipment (FE), and Safety and Performance Improvement Activities (SPI). Each domain includes specific items that gauge awareness of the standards and their implications for nursing practice. The scoring system typically employs a Likert scale format, where respondents rate their level of awareness from "Not at All Aware" to "Very Aware." Each response is assigned a numerical value (e.g., 1 for "Not at All Aware" to 5 for "Very Aware"), allowing for a cumulative score that reflects the overall level of awareness regarding nursing services accreditation standards. Higher scores indicate a greater familiarity with the standards, providing valuable insights for identifying areas needing further education and support.

### Data Analysis

#### Normality assumption

A normal data distribution is indicated by Z-value skewness and Z-value kurtosis falling within  $\pm 3$  and  $\pm 7$ , respectively. The study followed Hair et al. (2011) criteria, using a critical cut-off value of  $\pm 2.58$  [20]. The kurtosis and skewness values for each construct were observed to be within this range (Table 2). The descriptive analysis revealed an almost normal distribution, with mean skewness from -0.005 to 0.441 and kurtosis values between -0.864 and 0.058. Detailed skewness and kurtosis values can be found in Table 2.

**Table (2):** Skewness and Kurtosis for Variables.

Variables	Code	Skewness	Kurtosis
Awareness Regarding Accreditation	ARA	-.056	-.718
Organisation and Management	OM	.026	-.620
Human Resource Development	HRD	.438	.058
Policies and Procedures	PP	.441	-.507
Facilities and Equipment	FE	.302	-.624
Safety and Performance Improvement	SPI	-.005	-.864

## RESULTS AND DISCUSSION

### Assessment of confirmatory factor analysis (CFA) results (measurement model)


This study aimed to evaluate HCPs' awareness regarding accreditation through a structured analysis. An exploratory factor analysis (EFA) initially identified nine distinct factors related to accreditation awareness.

Following the EFA, a CFA integrated these factors into a cohesive model, including the ARA factor and five constructs based on MSQH standards: OM, HRD, PP, FE, and SPI. The integration allowed a thorough assessment of convergent and discriminant validity. To evaluate the model's fit, various fit indices were analyzed. The results indicated strong validity and reliability, evidenced by acceptable fit indices such as chi-squared over degrees of freedom (Chi-square/df), Root Mean Square Error of Approximation (RMSEA), and Comparative Fit Index (CFI). Specifically, Chi-square/df was found to be 1.982, CFI was 0.915, and RMSEA was 0.050, demonstrating a good fit according to established criteria.

### The Fitness of a measurement model

Brown (2015) recommends reporting fit indices such as chi-squared goodness of fit ( $\chi^2$ ), standardized root mean square residual (SRMR) for absolute fit, RMSEA for parsimony-corrected fit, and Tucker–Lewis fit index (TLI) and comparative fit index (CFI) for comparative fit. Mohamad, Awang, and Salim (2017) advised using at least one index from each of the three fitness categories: absolute fit, incremental fit, and parsimonious fit [21].

The present study employed commonly used fitness indices in scholarly research: chi-squared over degree of freedom (Chi-square/df), Root Mean Square Error of Approximation (RMSEA), and Comparative Fit Index (CFI) [22]. The model fit was determined based on specific cut-off values: RMSEA values between 0.05 and 1.00, CFI  $\geq$  0.90, and Chi-square/df  $\leq$  5.00, indicating a reasonable fit as per the criteria set by Hair et al. (2014) [22]. The results indicated that Chi-square/df = 1.982 ( $\leq$  5.00), CFI = .915, IFI = .916, TLI = .905 ( $\geq$  0.90), GFI = 0.859 ( $\leq$  0.85), RMSEA = .050 ( $\leq$  .080), and RMR = .026 ( $\leq$  0.10), which is considered a good fit, as shown in Figure 1 and Table 3.



**Figure (1):** Confirmatory Factor Analysis (CFA) Results for All Variables.

**Table (3):** Results of fit indices for CFA- Final Measurements Model.

Measure	Estimate	Threshold*	Interpretation
CMIN	1308.436	--	--
DF	660	--	--
CMIN/DF	1.982	<3	Excellent
CFI	0.915	>0.95	Acceptable
GFI	0.865	>0.85	Acceptable
IFI	0.916	>0.95	Acceptable
TLI	0.905	>0.95	Acceptable
RMSEA	0.050	<0.05	Excellent
RMR	0.026	<0.10	Excellent
P-value	0.000	>0.05	Acceptable

Source: Computed Data Analysis, \*(Awang, 2014; Hair et al. 2010)

### Reliability (Cronbach's Alpha) and composite reliability (CR)

The assessment of reliability ensures internal consistency when evaluating variables (Kim & Cha, 2002). A reliability score

above 0.60 is considered acceptable for the instrument [22,23]. This study conducted two types of reliability tests: Cronbach's alpha using SPSS 25.0 and CR. Specifically, the study found Cronbach's alpha values ranging from 0.751 for ARA.1 to 0.881 for SPI, while CR values ranged between 0.768 and 0.878 for the same variables as presented in Table 4. All the reliability and composite reliability values for the constructs reached a higher than the recommended threshold of 0.60.

#### Constructs validity

Convergent validity, the first type, was assessed through structural equation modelling (SEM) analysis in the measurement model to ascertain whether the indicators in a scale load together on a single construct. Discriminant validity, the second type, was used to confirm whether the items designed to measure different constructs are indeed evaluating different constructs

#### Convergent validity

Convergent validity assesses the degree to which measures of a specific construct align, indicating a high proportion of shared variance [22]. Convergent validity can be evaluated through CFA. According to Table 4, all items exhibited loadings above 0.50 on their respective constructs. A factor loading exceeding 0.50 is considered acceptable for a study sample of more than 300 respondents and provides strong evidence of convergent validity [20]. With factor loadings ranging from (0.552) for HRD13 to (0.852) for OM2, all indicators in the study were clearly associated with their respective constructs, providing solid support for the model's convergent validity.

Furthermore, Table 4 shows the Average Variance Extracted (AVE) test results for two constructs [human resource development1 (HRD.1) and awareness regarding accreditation2 (ARA.2)], which were below the recommended threshold of 0.5 (0.451 and 0.455, respectively). Consistent with previous research [24], while an AVE exceeding 0.5 is preferred, values as low as 0.4 are acceptable. Following the criteria set by Fornell and Larcker (1981), an AVE below 0.5, but with a composite reliability (CR) greater than 0.6, indicates adequate convergent validity of the construct [25]. Table 4.19 presents the findings of item loadings, AVE, Cronbach's alpha, and CR.

**Table (4):** Items loading, Cronbach's alpha, Composite Reliability (CR) and (AVE).

Constructs	Items	Factor Loading	Cronbach's alpha	CR	AVE
ARA.1	ARA3	0.622	0.751	0.768	0.455
	ARA4	0.592			
	ARA5	0.731			
	ARA6	0.641			
	ARA9	0.675			
ARA.2	ARA10	0.772	0.858	0.868	0.570
	ARA11	0.823			
	ARA12	0.751			
	ARA13	0.746			
OM.1	OM1	0.713	0.807	0.811	0.591
	OM2	0.852			
	OM3	0.733			
OM.2	OM5	0.653	0.851	0.856	0.545
	OM6	0.796			
	OM7	0.806			
	OM8	0.742			
	OM9	0.681			
HRD.1	HRD6	0.777	0.769	0.760	0.516
	HRD7	0.724			
	HRD9	0.647			
HRD.2	HRD10	0.743	0.786	0.758	0.451
	HRD11	0.746			
	HRD12	0.603			
	HRD13	0.552			
PP	PP5	0.653	0.858	0.854	0.543
	PP6	0.667			
	PP7	0.789			

Constructs	Items	Factor Loading	Cronbach's alpha	CR	AVE
FE	PP8	0.798	0.861	0.857	0.548
	PP9	0.817			
	FE1	0.622			
	FE2	0.732			
	FE3	0.821			
	FE4	0.762			
SPI	FE5	0.749	0.881	0.878	0.591
	SPI2	0.751			
	SPI3	0.842			
	SPI4	0.734			
	SPI5	0.780			
	SPI6	0.732			

Source: Prepared by researcher using output of Amos-SEM (Measurement Model).

#### Discriminant validity

Discriminant validity refers to the extent to which a particular construct is distinct from other constructs [20]. It is assessed by comparing the average variance extracted (AVE) for each construct with the squared correlations between that construct and others [25]. In our analysis, the AVE values for each construct exceeded the squared correlations with other constructs, demonstrating discriminant validity. Specifically, the square root of the AVE for each construct was greater than the absolute value of the correlation squared with any other factor ( $AVE > \text{correlation square}$ ). This was further supported by Table 5, which shows that the AVE and the square root of the AVE for all constructs exceeded the correlations with other constructs in the model.

**Table (5):** Discriminant Validity for Latent Variables for Fornell & Larcker, (1981).

HRD.1	FE	OM.1	HRD.2	OM.2	SPI	ARA.2	ARA.1	PP
0.718								
0.527	0.740							
0.515	0.456	0.768						
0.712	0.458	0.462	0.672					
0.493	0.537	0.571	0.447	0.738				
0.588	0.663	0.507	0.525	0.616	0.769			
0.248	0.286	0.210	0.174	0.378	0.276	0.755		
0.295	0.353	0.283	0.308	0.365	0.393	0.462	0.675	
0.498	0.640	0.388	0.532	0.513	0.602	0.276	0.309	0.737

Source: Prepared by researcher using output of Amos-SEM (Measurement Model).

Note: HRD.2=Human Resource Development2 HRD.1= Human Resource Development1; FE= Facilities and Equipment; OM.2= Organization and Management2; OM.1= Organization and Management1; SPI= Safety and Performance Improvement; ARA.2= Awareness Regarding Accreditation; ARA.1= Awareness Regarding Accreditation1; PP= Policies and Procedure.

## DISCUSSION

This study successfully developed and validated a measurement model to assess the awareness and challenges surrounding nursing services accreditation among healthcare providers (HCPs) in Yemen. By employing rigorous statistical methods, including Confirmatory Factor Analysis (CFA), the research established the construct validity, reliability, and overall robustness of nine distinct constructs: Awareness Regarding Accreditation (ARA), Organization and Management (OM), Human Resource Development (HRD), Policies and Procedures (PP), Facilities and Equipment (FE), and Safety and Performance Improvement (SPI). The use of well-established fit indices confirmed that the model demonstrated adequate fitness, thereby validating the effectiveness of the measurement instrument.

The identified constructs are essential for understanding the current landscape of nursing services accreditation in Yemen. For instance, the strong emphasis on organizational management and human resource development suggests that healthcare institutions need to enhance their administrative

frameworks and invest in ongoing training for staff [3, 26, 27]. These factors are crucial for fostering an environment conducive to high-quality patient care, especially in resource-limited settings where healthcare infrastructure may be underdeveloped [28,29].

Notably, the results revealed that while the measurement model was robust, some constructs exhibited Average Variance Extracted (AVE) values slightly below the preferred threshold of 0.5. Specifically, constructs related to Human Resource Development (HRD) and Awareness Regarding Accreditation (ARA) had AVE values of 0.451 and 0.455, respectively. While these values suggest room for improvement, they still meet the criteria for adequate convergent validity when accompanied by satisfactory composite reliability scores. This indicates that the

instruments can still effectively capture the constructs they are intended to measure, although further refinement may enhance their validity [31,32].

The practical implications of these findings are significant. The validated instrument serves as a critical tool for identifying specific areas requiring attention in the accreditation process, such as the development of comprehensive training programs. It emphasizes the importance of continuous education not only for HCPs but also for healthcare managers and policymakers [33, 34]. Cultivating a culture of quality and accreditation is essential, particularly in regions like Hadhramaut, where gaps in knowledge and resistance to change can impede progress. Engaging HCPs in regular training initiatives can significantly enhance their understanding of the accreditation process and its role in improving patient outcomes and healthcare performance [3, 35].

Moreover, this study highlighted the need for future research to address the limitations of the current sample, which was restricted to doctors, nurses, and managers. Incorporating a broader range of healthcare professionals, such as laboratory technicians, pharmacists, and support staff, would provide a more comprehensive perspective on the challenges and opportunities associated with nursing services accreditation. By expanding the scope of future studies, researchers can capture a fuller understanding of the accreditation landscape and develop strategies that encompass the insights of all relevant stakeholders.

## CONCLUSION

This study explored the relationships among Gacha game use, sleep quality, and aggression among university students. The findings indicate that Gacha game players experience poorer sleep quality and moderate levels of aggression. However, no significant correlations were found between game duration, sleep quality, and aggression, possibly due to cultural factors or the catharsis theory. These results highlight the need for increased awareness of the potential negative consequences of excessive Gacha game use and the importance of monitoring and self-regulation to minimize adverse effects.

## Disclosure Statement

- **Ethics approval and consent to participate:** The Institutional Review Board Hadhramaut University approved this study on 20/6/2023 (Ref no. H.U.1/16/2023-2024). The current study followed the Declaration of Helsinki provisions, and all participants provided informed consent. The permission to use the measurement instruments was received from the original author.
- **Consent for publication:** The authors grant the Publisher permission to publish this work. All the data generated for this study are included within the article.
- **Availability of data and materials:** All data generated

during this study are included in this published article.

- **Conflicts of interest:** The authors report no conflict of interest.
- **Author's contribution;** study conception and design: Talal Mansoor, Sharifa Ezat Bint Wan Puteh, Azimatun Noor Aizuddin, Nawal Saeed Banafa; data analysis and validation, Talal Mansoor, Malakeh. Z. Malak, Sharifa Ezat Bint Wan Puteh; draft manuscript preparation: Talal Mansoor, Malakeh. Z. Malak, Sharifa Ezat Bint Wan Puteh, Azimatun Noor Aizuddin, Nawal Saeed Banafan. All authors reviewed the results and approved the final version of the manuscript.
- **Funding:** None.
- **Acknowledgements:** The authors would like to thank all nurses who participate in the study and hospital administrators who facilitate the data collection process.

## Open Access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <https://creativecommons.org/licenses/by-nc/4.0/>

## References

- 1] Ružic-Baf, M., H. Strnak, and A. Debeljuh, Online Video Games and Young People. *International Journal of Research in Education and Science*, 2016. 2(1): p. 94-103.
- 2] Gealan, R. 2015. NURSES' EXPERIENCE ON INTERNATIONAL ACCREDITATION. The Philippine Women's University.
- 3] Jun, J., Ojemeni, M. M., Kalamani, R., Tong, J., & Crecelius, M. L. (2021). Relationship between nurse burnout, patient and organizational outcomes: Systematic review. *International journal of nursing studies*, 119, 103933
- 4] Alnaeem, M. M., Hamdan-Mansour, A. M., Nashwan, A. J., Abuataallah, A., & Al-Hussami, M. (2022). Healthcare providers' intention to leave their jobs during COVID-19 pandemic: A cross-sectional study. *Health Science Reports*, 5(6), e859
- 5] Waddill-Goad, S. (2023). Beyond Burnout: Overcoming Stress in Nursing & Healthcare for Optimal Health & Well-being. Sigma Theta Tau
- 6] Katoue, M. G., Somerville, S. G., Barake, R., & Scott, M. (2021). The perceptions of healthcare professionals about accreditation and its impact on quality of healthcare in Kuwait: a qualitative study. *Journal of evaluation in clinical practice*, 27(6), 1310-1320
- 7] Mansour, W., Boyd, A., & Walshe, K. (2020). The development of hospital accreditation in low-and middle-income countries: a literature review. *Health policy and planning*, 35(6), 684-700
- 8] Wardhani, V., Van Dijk, J.P. & Utarini, A. 2019. Hospitals accreditation status in Indonesia: Associated with hospital characteristics, market competition intensity, and hospital performance? *BMC Health Services Research* 19(1).



- 9] Hussein, M., Pavlova, M., Ghalwash, M., & Groot, W. (2021). The impact of hospital accreditation on the quality of healthcare: a systematic literature review. *BMC health services research*, 21, 1-12
- 10] Molski, A. 2020. CREATING A FOUNDATION OF QUALITY IMPROVEMENT THROUGH ACCREDITATION STRATEGIES.
- 11] World Health Organization. 2016. Health System Profile-Yemen. Regional Health System Observatory. Report No.
- 12] Dureab, F., Hussain, T., Sheikh, R., Al-Dheeb, N., Al-Awlaqi, S., & Jahn, A. (2021). Forms of health system fragmentation during conflict: the case of Yemen. *Frontiers in Public Health*, 9, 659980
- 13] Al Waziza, R., Sheikh, R., Ahmed, I., Al-Masbhi, G., & Dureab, F. (2023). Analyzing Yemen's health system at the governorate level amid the ongoing conflict: a case of Al Hodeida governorate. *Discover Health Systems*, 2(1), 15
- 14] Abbad, I. B. (2024). Challenges of Local Public Administration Dministrtrtion in Conflict-Affected Countries: Case of Hadhramaut Governorate in Yemen
- 15] Ali, E. A. S. (2020). Providing Health Care in Armed Conflict Countries: The Case of Yemen (Master's thesis, Hamad Bin Khalifa University (Qatar))
- 16] Nasser, A. A., Al-Ashwal, M. M., Al-Khulaidi, A. A., Al-Sana'ani, S. S., & Al-jobber, M. Assessing Equity in Healthcare Facility Resource Allocation in Yemen: An Entropy-TOPSIS Analysis
- 17] Ba-Break, M., Donnelly, S., Saleh, M., Kaity, A. M., Alwaqedi, A., Badheeb, A. M., ... & Annuzaili, D. A. (2024). COVID-19 surveillance in fragile health systems, armed conflict and humanitarian crisis, the case of Yemen. *International Journal of Healthcare Management*, 1- 13
- 18] Mansoor, T., Puteh, S. E. W., Aizuddin, A. N., & Malak, M. Z. (2024). Challenges and Strategies in Implementing Hospital Accreditation Standards Among Healthcare Professionals in Healthcare Systems in Yemen: A Phenomenological Study. *Cureus*, 16(4)
- 19] Alraimi, A. A., & Al-Nashmi, M. M. (2024). The interactive effect of applying the management-centered standards of Joint Commission International (JCI) and practicing administrative control in improving the quality of health services: a study on three Yemeni hospitals seeking accreditation. *Journal of Hospital Management and Health Policy*, 8
- 20] Badr, A., & Abol-Ghaith, F. M. (2024). Health workforce in governmental hospitals Hajjah Governorate-Yemen 2022
- 21] Hair, J.F., Ringle, C.M. & Sarstedt, M. 2011. PLS-SEM: Indeed a silver bullet. *Journal of Marketing Theory and Practice* 19(2): 139–152.
- 22] Mohamad, M.; Awang, Z.; Salim, N.A.M.A. 2017. Validating the Maqasid Shariah Prison Quality of Life (MSPQoL) among drug-abuse inmates using confirmatory factor analysis. *Int. J. Appl. Bus. Econ. Res.* 2017, 15, 91–103.
- 23] Hair, J. F., Gabriel, M., & Patel, V. (2014). AMOS covariance-based structural equation modeling (CB-SEM): Guidelines on its application as a marketing research tool. *Brazilian Journal of Marketing*, 13(2).
- 24] Sekaran, S., Foster, R. G., Lucas, R. J., & Hankins, M. W. (2003). Calcium imaging reveals a network of intrinsically light-sensitive inner-retinal neurons. *Current biology*, 13(15), 1290-1298
- 25] Barclay, D., Higgins, C., & Thompson, R. (1995). The partial least squares (PLS) approach to casual modeling: personal computer adoption ans use as an Illustration.
- 26] Fornell, C., & Larcker, D. F. (1981). Evaluating structural equation model with unobservable variables and measurement error. *Journal of Marketing Research* 18 (1), 39-50.
- 27] Labrague, L. J., Al Sabei, S., AbuAlRub, R., Burney, I., & Al Rawajfah, O. (2024). The role of nurses' adherence to clinical safety guidelines in linking nurse practice environment to missed nursing care. *Journal of Nursing Scholarship*
- 28] Alzoubi, M. M., Al-Mugheed, K., Oweidat, I., Alrahbeni, T., Alnaeem, M. M., Alabdullah, A. A. S., ... & Hendy, A. (2024). Moderating role of relationships between workloads, job burnout, turnover intention, and healthcare quality among nurses. *BMC psychology*, 12(1), 1-9
- 29] Soósová, M. S. (2021). Association between nurses' burnout, hospital patient safety climate and quality of nursing care. *Central European Journal of Nursing and Midwifery*, 12(1), 245
- 30] Labrague, L. J. (2022). Linking nurse practice environment, safety climate and job dimensions to missed nursing care. *International nursing review*, 69(3), 350-358
- 31] Suleiman, K., Al Kalaldehy, M., AbuSharour, L., Yates, B., Berger, A., Mendoza, T., ... & Menshawi, A. (2019). Validation study of the Arabic version of the Brief Fatigue Inventory (BFI-A). *Eastern Mediterranean Health Journal*, 25(11), 784-790
- 32] Suleiman, K., Al Kalaldehy, M., AbuSharour, L., Yates, B., Berger, A., Mendoza, T., ... & Menshawi, A. (2019). Validation study of the Arabic version of the Brief Fatigue Inventory (BFI-A). *Eastern Mediterranean Health Journal*, 25(11), 784-790
- 33] Clark, L. A., & Watson, D. (2019). Constructing validity: New developments in creating objective measuring instruments. *Psychological assessment*, 31(12), 1412
- 34] Garcia, C. D. L., Abreu, L. C. D., Ramos, J. L. S., Castro, C. F. D. D., Smiderle, F. R. N., Santos, J. A. D., & Bezerra, I. M. P. (2019). Influence of burnout on patient safety: systematic review and meta- analysis. *Medicina*, 55(9), 553
- 35] Alnaeem, M. M., Sabra, M. A. A., Jebbeh, R. A., & Suleiman, K. (2024). Workplace violence against nurses in rural governmental hospitals in Jordan. *Collegian*, 31(5), 348-355
- 36] Alshamsi, A. I., Thomson, L., & Santos, A. (2020). What impact does accreditation have on workplaces? A qualitative study to explore the perceptions of healthcare professionals about the process of accreditation. *Frontiers in Psychology*, 11, 1614