



# Solving Higher Order Ordinary Differential Equations in Physics: A New Modified Adomian Decomposition Method Approach to MHD Flows, Elastic Beam, and Sixth-Order Boundary value problem

Received 13 April 2025, Accepted 16 May , 2025, Published xxxx, DOI: <https://doi.org/10.xxxx>

Nuha Mohammed Dabwan<sup>1</sup>, Yahya Qaid Hasan<sup>2</sup>,

**Abstract:** The study presents a novel approach for solving high-order ordinary differential equations (ODEs) prevalent in various physics applications, specifically through a modified Adomian Decomposition Method (ADM). This method enhances the traditional ADM by introducing specific modifications that improve its convergence and applicability to complex problems. The research focuses on three primary areas: magnetohydrodynamic (MHD) flows, the dynamics of elastic beams, and sixth-order boundary value problems. The proposed method demonstrates significant effectiveness in deriving analytical solutions that can accurately predict physical behaviors in these domains. By applying the modified ADM, the study not only addresses the challenges associated with higher order ODEs but also offers practical solutions for engineers and physicists working with intricate modeling scenarios. The results indicate that this method provides an efficient and reliable framework for analyzing and solving complex differential equations in the field of physics and engineering.

**Keywords:** Magnetohydrodynamic (MHD) Flows; Jeffery-Hamel Problem; Elastic Beam Equation; Sixth-Order Boundary Value Problem; Modified Adomian Decomposition Method; Boundary Conditions.

## Introduction

Differential equations serve as a fundamental tool across various scientific disciplines, including physics, chemistry, and engineering. They provide a framework for modeling an extensive range of natural phenomena, allowing for a deeper understanding of dynamic systems characterized by either linear or nonlinear behavior. By developing models based on these equations, researchers can derive both analytical and approximate solutions, which have gained significant prominence in recent studies.

Given the inherent challenges in obtaining precise solutions for physical problems represented mathematically by differential equations, the exploration of analytical and numerical techniques becomes imperative. Foundational work in operator calculus, such as that of Mikusiński and Riemann–Liouville [1–3], has provided a theoretical basis for manipulating and inverting differential operators, which underpins many decomposition-based methods.

In fluid mechanics, for instance, the Jeffery-Hamel flow, a significant conceptual advancement introduced by Jeffery [4] and Hamel [5] in the early 20th century, has become a pivotal example of how two-dimensional incompressible viscous flows can be examined using Navier-Stokes equations.

The term magnetohydrodynamic (MHD) was first introduced by re-

searcher Parth Bansal, marking a significant development in the field of fluid mechanics [6]. MHD studies the behavior of electrically conducting fluids in the presence of magnetic fields, combining principles from both electromagnetism and fluid dynamics. Bansal's contributions laid the groundwork for exploring various MHD applications, particularly in contexts like astrophysics, engineering, and plasma physics, where magnetic fields interact with fluid flow, influencing behaviors such as turbulence and stability [7].

The diverse physical problems associated with fourth-order differential equations are intrinsically linked to the field of elastic stability theory. These equations play a critical role in modeling the behavior of structural elements under various loading conditions, particularly those that exhibit bending and torsional deformations. When addressing the elastic stability of structures, one encounters phenomena such as buckling, which occurs when a structure is subjected to critical loads that exceed its capacity to maintain its original shape [8]. The analysis of such problems typically involves fourth-order differential equations due to their ability to incorporate the effects of shear deformation and rotational inertia, which are essential for accurately predicting the stability and performance of structural components [9–12].

A comprehensive and uniform framework for sixth-order boundary value problems is increasingly recognized as a vital tool for analyzing a diverse array of challenges across fields such as astrophysics, mathe-

<sup>1</sup>Department of Mathematics

Faculty of Education-Aden, University of Aden, Aden, Yemen

Faculty of Education-AL-Mekhlaf, Taiz University, Taiz, Yemen

E-mail: [nuhamdm@gmail.com](mailto:nuhamdm@gmail.com)

<sup>2</sup>Department of Mathematics

Faculty of Education and Sciences, University of Saba Region, Marib, Yemen

E-mail: [qaid.yahya@tu.edu.ye](mailto:qaid.yahya@tu.edu.ye)

matics, and engineering sciences. This framework allows researchers and practitioners to study complex systems and phenomena that exhibit higher-order behavior, which is often crucial for accurately modeling the intricate processes in these domains [13].

In astrophysics, for instance, many phenomena, including the convective stability in stellar structures or the dynamics of astronomical bodies, can be effectively described using sixth-order differential equations. These equations arise from the need to account for higher-order effects that are significant in understanding stability and oscillatory behaviors, allowing for a more nuanced description of astrophysical systems [14]. Given the inherent challenges in obtaining precise solutions for physical problems represented mathematically by differential equations, the exploration of numerical techniques becomes imperative. Among these, the Adomian Decomposition Method (ADM) [15, 16] has demonstrated its utility, particularly in addressing ordinary differential equations without the need for discretization or linearization, and its performance has been enhanced by a number of modifications [17–27]. ADM provides an approximate solution of the problem without simplifying it, in contrast to usual methods that include mild nonlinearity and minor perturbations, which alter the physics of the problem through simplification.

The research conducted by various authors illustrates the versatility of the (ADM) and its modified versions in tackling complex differential equations across multiple domains. Odiba's work exemplifies the application of the Modified ADM (MADM) to a third-order ordinary differential equation, showcasing its efficacy in deriving solutions that might be computationally intensive with traditional methods [28]. Meanwhile, I. Hashim's integration of ADM for a fourth-order integro-differential equation highlights the method's adaptability in handling intricate mathematical models, such as those found in engineering and physics [29]. While Ali et al.'s application of ADM to the HIV infection model indicates its relevance in biological systems modeling [30]. Similarly, Coskun et al. demonstrated ADM's utility in studying the behavior of Euler beams under variable stiffness, reinforcing the method's significance in structural analysis [31].

However, most existing operator-based extensions of the Adomian Decomposition Method (ADM) focus on polynomial or standard differential operator forms and do not explicitly address exponential-type differential operators. This highlights a gap in the current literature, which our proposed modification seeks to fill by incorporating complex exponential weightings into the operator structure. This study aims to develop and demonstrate a modified Adomian Decomposition Method (ADM) for addressing higher order ordinary differential equations (ODEs) relevant to physics. The focus is on applying this method to specific problems, including magnetohydrodynamic (MHD) flows, the behavior of elastic beams, and sixth-order boundary value problems.

The goal of the research is to provide an effective analytical solution technique that enhances the flexibility and applicability in solving complex ODEs. This approach is intended to bridge theoretical advancements and practical applications in fields such as physics and engineering, ultimately aiming to offer more robust solutions to problems characterized by higher order differential equations in various physical contexts.

## The Mathematical Formulation of the Jeffery-Hamel flow and the Beam Equation of Fourth Order

### First

As shown in Figure 1, the Figure represent the steady two-dimensional flow of an incompressible, conductive, viscous fluid between two rigid plane walls that meet at an angle of  $2\alpha$ . The velocity of the fluid depends solely on the radial distance ( $r$ ) and the angle ( $\theta$ ), and it is entirely radial in nature [32]. In polar coordinates, the continuity equation and the Navier-Stokes equations can be expressed as follows:

$$\frac{\rho}{r} \frac{\partial}{\partial r} (ru(r, \theta)) = 0, \quad (1)$$

$$u(r, \theta) \frac{\partial u(r, \theta)}{\partial r} + \frac{1}{\rho} \frac{\partial p}{\partial r} + \frac{\sigma B_0^2}{\rho r^2} u(r, \theta) - \mu \left[ \frac{\partial^2 u(r, \theta)}{\partial r^2} + \frac{1}{r} \frac{\partial u(r, \theta)}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u(r, \theta)}{\partial \theta^2} - \frac{u(r, \theta)}{r^2} \right] = 0, \quad (2)$$

$$\frac{1}{\rho r} \frac{\partial p}{\partial \theta} - \frac{2\mu}{r^2} \frac{\partial u(r, \theta)}{\partial \theta} = 0, \quad (3)$$

note that,  $\rho$  the fluid density,  $\mu$  is the coefficient of kinematic viscosity,  $\sigma$  the conductivity,  $B_0$  the electromagnetic induction and  $p$  is the fluid pressure.

Using equation (1)

$$h(\theta) = ru(r, \theta), \quad (4)$$

from dimensionless parameters

$$f(x) = \frac{h(\theta)}{h_{max}}, \quad x = \frac{\theta}{\alpha}, \quad (5)$$

Putting them in (2) and (3), we get nonlinear third order boundary value problem

$$f'''(x) + 2\alpha Re f(x) f'(x) + (4 - Ha) \alpha^2 f'(x) = 0, \quad (6)$$

with boundary conditions

$$f(0) = 1, f'(0) = 0, f(1) = 0. \quad (7)$$

Where  $Re$  is Reynolds number

$$Re = \frac{U_{max}}{\mu} \begin{pmatrix} \text{divergent} : \alpha > 0, U_{max} > 0 \\ \text{convergent} : \alpha < 0, U_{max} < 0 \end{pmatrix},$$

and  $Ha$  is Hartmann number

$$Ha = \sqrt{\frac{\sigma B_0^2}{\rho \mu}}.$$




Figure 1: Geometry of the MHD Jeffery-Hamel flow.

### Second

When a beam-column that subjected to both the axial load and the spread load perpendicular to axis as indicated in figure 2 (b) has a cross section distance  $dx$  internal forces occur in the element in figure 2 (a). When the equilibrium equation is expressed in the direction of the distributed  $q$ , it is given as:

$$q = -\frac{dV}{dx}. \quad (8)$$

The equilibrium equation in the  $y$  direction can be written as an ordinary differential equation, as presented in [33–35]

$$-V + qdx + (V + dV) = 0. \quad (9)$$

The sum of the forces acting on each surface of the cross-section must be balanced due to the equilibrium condition. This can be expressed as:

$$M + qdx + \frac{dx}{q} + (V + dV)dx - (M + dM) + p \frac{dy}{dx} = 0. \quad (10)$$

Here  $M$  is the bending moment trying to bend the cross-section element, and  $V$  is the shear force acting on the surface of the element. Assuming the rotation is small and neglecting second order terms in  $dx$ , equation (10) simplifies to:

$$V = \frac{dM}{dx} - p \frac{dy}{dx}. \quad (11)$$

Since small rotation is assumed, and considering that  $\frac{d^2}{dx^2} = -\frac{M}{EI}$ , then (11) written as

$$-V = EI \frac{d^3 y}{dx^3} + p \frac{dy}{dx}. \quad (12)$$

Here,  $EI$  refers to bending rigidity. Differentiating both sides of equation (12) with respect to  $x$ , we obtain a fourth-order differential equation for the elastic curve, given by:

$$EI \frac{d^4 y}{dx^4} + p \frac{d^2 y}{dx^2} = q(x). \quad (13)$$




Figure 2: Cross-section analysis of the column-beam element.

## New Modified ADM

In this section, we present an innovative modification to the Standard Adomian Decomposition Method (ADM), specifically focusing on alterations made to the differential operator. This enhancement aims to improve the method's applicability and effectiveness in solving nonlinear differential equations.

Consider the following equation:

$$y^{(r+2)} + \nu y^{(r)} = g(x) + N(y), \quad r \in N \cup \{0\}, \quad (14)$$

with the conditions

$$y(0) = a_0, y'(0) = a_1, \dots, y^{(r+1)}(0) = a_i, i = 0, 1, \dots,$$

in the standard Adomian Decomposition Method (SADM), the linear operator is typically taken as  $Ly = y^{(r+2)}$ , where  $L(y) = y^{(r+2)}$ . However, in our proposed modification, we define the linear operator as  $Ly = y^{(r+2)} + \nu y^{(r)}$ , where  $r = n + m$ .

The choice  $Ly = y^{(r+2)} + \nu y^{(r)}$  improves convergence because it modifies the spectral characteristics of the linear operator, leading to a more balanced and well-conditioned inverse. This structure helps reduce the growth of higher-order terms in the decomposition series, thereby accelerating convergence, especially for stiff or oscillatory

problems.

Where  $L(y)$  as the following

$$L_1(y) = \frac{d^m}{dx^m} e^{\pm i \sqrt{\nu} x} \frac{d}{dx} e^{\mp 2i \sqrt{\nu} x} \frac{d}{dx} e^{\pm i \sqrt{\nu} x} \frac{d^n}{dx^n} (y), \quad (15)$$

$$L_2(y) = \frac{d^m}{dx^m} e^{\pm i \sqrt{\nu} x} \frac{d}{dx} e^{\mp i \sqrt{\nu} x} \frac{d^n}{dx^n} e^{\mp i \sqrt{\nu} x} \frac{d}{dx} e^{\pm i \sqrt{\nu} x} (y). \quad (16)$$

Where  $\nu$  is constant and in equation (15)  $n, m \in N \cup \{0\}$ , and in equation (16)  $n \in N$  and  $m \in N \cup \{0\}$ .

The inverse of eq.(15) and eq.(16) as follows

$$L_1^{-1}(y) = \underbrace{\int_0^x \int_0^x \dots \int_0^x}_{n\text{-times}} e^{\mp i \sqrt{\nu} x} \int_0^x e^{\pm 2i \sqrt{\nu} x} \int_0^x e^{\mp i \sqrt{\nu} x} \underbrace{\int_0^x \int_0^x \dots \int_0^x}_{m\text{-times}} (y) dx dx \dots dx dx dx dx \dots dx. \quad (17)$$

$$L_2^{-1}(y) = e^{\mp i \sqrt{\nu} x} \int_0^x e^{\pm i \sqrt{\nu} x} \underbrace{\int_0^x \int_0^x \dots \int_0^x}_{n\text{-times}} e^{\pm i \sqrt{\nu} x} \int_0^x e^{\mp i \sqrt{\nu} x} \underbrace{\int_0^x \int_0^x \dots \int_0^x}_{m\text{-times}} (y) dx dx \dots dx dx dx dx \dots dx. \quad (18)$$

So equation (14) takes the form

$$Ly + Ny = g(x),$$

or equivalently:

$$Ly = g(x) - Ny, \quad (19)$$

to get  $y$ , take  $L^{-1}$  ( $L_1^{-1}$  or  $L_2^{-1}$ ) to both sides of equations (19). Then

$$y = \kappa(x) + L^{-1}g(x) - L^{-1}N(y), \quad (20)$$

where  $\kappa(x)$  arising from auxiliary conditions.

The solution in ADM decomposition to [15]

$$y_0 = \kappa(x) + L^{-1}g(x),$$

and

$$y_{n+1} = -L^{-1}N(y),$$

where

$$N(y) = \sum_{n=0}^{\infty} A_n,$$

$A_n$  called Adomian polynomial and formed by

$$A_n = \frac{1}{n!} \frac{d^n}{d\lambda^n} \left[ N \left( \sum_{i=0}^{\infty} \lambda^i y_i \right) \right], \quad \text{where } \lambda = 0, n = 0, 1, 2, 3, \dots \quad (21)$$

That is

$$y_{n+1} = -L^{-1}A_n,$$

then

$$y_1 = -L^{-1}A_0,$$

$$y_2 = -L^{-1}A_1,$$

$$y_3 = -L^{-1}A_2,$$

and so on.

Based on the above, the solution via ADM is given as a series:

$$y(x) = y_0 + y_1 + y_2 + y_3 + \dots = \sum_{n=0}^{\infty} y_n. \quad (22)$$

To better illustrate how the proposed NMADM differs from other modified ADM approaches, Table 1 provides a comparative summary of the key features, convergence strategies, and implementation considerations.

## Justification for Using Complex Exponential Weightings

The composition of differential operators with complex exponential weightings in equations (15) and (16) is motivated by their ability to reshape the operator's structure and improve its spectral properties. Exponential terms like  $e^{\mp 2i\sqrt{\nu}x}$  act as integrating factors or transformation kernels, which can reduce oscillatory behavior and enhance the stability of the inverse operator. This strategy is inspired by techniques in spectral methods and perturbation theory, where such weightings are commonly used to simplify the inversion of differential operators and accelerate convergence. As a result, the modified operator becomes better suited for handling nonlinearities and leads to more efficient implementation of the decomposition process.

## Existence and Uniqueness of Solution

Let  $\aleph : \hbar \rightarrow \hbar$  is an operator, where  $\hbar$  is the Banach space  $(C(I_0), \|\cdot\|)$ , the space of all continuous functions on  $I_0$  with norm  $\|y\| = \max_{x \in I_0} |y(x)|$ , and  $N$  satisfied Lipschitz condition  $(|N(y) - N(z)| \leq \ell|y - z| \text{ where } y, z \in \hbar)$ .

**Theorem 1** *The equation (14) has a unique solution  $y \in \hbar$  on  $I_0 = [0, T]$  when  $0 < \varrho < 1$  where  $\varrho = \ell\gamma$ , and  $\gamma = \sum_{i=0}^{\infty} (-1)^i \nu^i \frac{T^{(r+2)+2i}}{((r+2)+2i)!}$ .*

**proof** *The mapping  $\aleph : \hbar \rightarrow \hbar$  is defined as*

$$(\aleph y)(x) = \kappa(x) + L^{-1}g(x) - L^{-1}N(y),$$

$$(\aleph z)(x) = \kappa(x) + L^{-1}g(x) - L^{-1}N(z),$$

where  $L^{-1}$  define in equations (17,18).

$$\begin{aligned} \|(\aleph y)(x) - (\aleph z)(x)\| &= \max_{x \in I_0} |L^{-1}N(y) - L^{-1}N(z)| \\ &= \max_{x \in I_0} |L^{-1}(N(y) - N(z))| \\ &\leq \ell \max_{x \in I_0} |y - z| L^{-1}(1) \\ &\leq \ell\gamma \max_{x \in I_0} |y - z| \\ &\leq \varrho \|y - z\| \end{aligned}$$

Since  $0 < \varrho < 1$ , Then the mapping  $\aleph$  is contraction. That is, there exists a unique solution  $y \in \hbar$  by Banach contraction principle.

## Convergent and Error Estimate

In this section, we will study the Convergent and Error Estimate of the solution.

## Proof of Convergent

**Theorem 2** *The series solution (22) of the equation (14) using NMADM converges if  $|y_1| < \infty$  and  $0 < \varrho < 1$ ,  $\varrho = \ell\gamma$ .*

**Proof** Let  $S_n = \sum_{i=0}^n y_i$  is the sequence of partial sum. Since,

$$N(y) = N\left(\sum_{i=0}^{\infty} y_i\right) = \sum_{i=0}^{\infty} A_i,$$

so,

$$N(S_n) = \sum_{i=0}^n A_i.$$

Now, we prove that  $S_n$  is a Cauchy sequence in Banach space  $\hbar$ . Let  $S_n, S_m$  be two arbitrary partial sums such that  $n \geq m$ , then,

$$\begin{aligned} \|S_n - S_m\| &= \max_{x \in I_0} |S_n - S_m| = \max_{x \in I_0} \left| \sum_{i=m+1}^n y_i(x) \right| \\ &= \max_{x \in I_0} \left| \sum_{i=m+1}^n L^{-1} N y \right| \\ &= \max_{x \in I_0} \left| \sum_{i=m+1}^n L^{-1} A_i \right| \\ &= \max_{x \in I_0} \left| \sum_{i=m}^{n-1} L^{-1} A_i \right| \\ &= \max_{x \in I_0} |L^{-1}[N(S_n) - N(S_m)]| \\ &\leq \max_{x \in I_0} L^{-1} |N(S_n) - N(S_m)| \\ &\leq \max_{x \in I_0} L^{-1} \ell |S_n - S_m| \\ &\leq \gamma \ell \|S_n - S_m\| \\ &\leq \varrho \|S_n - S_m\|. \end{aligned}$$

Let  $n = m + 1$  then,

$$\begin{aligned} \|S_{m+1} - S_m\| &\leq \varrho \|S_m - S_{m-1}\| \\ &\leq \varrho^2 \|S_{m-1} - S_{m-2}\| \leq \dots \leq \varrho^m \|S_1 - S_0\|. \end{aligned}$$

Based on the triangle inequality, we get

$$\begin{aligned} \|S_n - S_m\| &\leq \|S_{m+1} - S_m\| + \\ &\|S_{m+2} - S_{m+1}\| + \dots + \|S_n - S_{n-1}\| \\ &\leq (\varrho^m + \varrho^{m+1} + \dots \varrho^{n-1}) \|S_1 - S_0\| \\ &\leq \varrho^m (1 + \varrho + \dots \varrho^{n-m-1}) \|S_1 - S_0\| \\ &\leq \varrho^m \left( \frac{1 - \varrho^{n-m}}{1 - \varrho} \right) \|y_1\|. \end{aligned}$$

Since  $0 < \varrho < 1$ ,  $(\varrho^{n-m}) \leq 1$ . Then

$$\|S_n - S_m\| \leq \frac{\varrho^m}{1 - \varrho} \|y_1\|$$

$$\leq \frac{\varrho^m}{1 - \varrho} \max_{x \in I_0} |y_1|.$$

Since  $|y_1| < \infty$ , then,  $\|S_n - S_m\| \rightarrow 0$  hence,  $S_n$  is a Cauchy sequence in Banach space  $\hbar$ , that is  $\sum_{n=0}^{\infty} y_n$  converges and this completes the proof.

## Error Estimate

The following theorem allows us to estimate the maximum absolute truncation error of the Adomian's series solution for NMADM.

**Theorem 3** The series solution (22) to the problem (14) is estimated to have a maximum absolute truncation error given by

$$\max_{x \in I_0} |y(x) - \sum_{i=0}^m y_i(x)| \leq \frac{\varrho^m}{1-\varrho} \max_{x \in I_0} |y_1|$$

**Proof** Theorem (2) gives us

$$\|S_n - S_m\| \leq \frac{\varrho^m}{1-\varrho} \max_{x \in I_0} |y_1|.$$

Since  $S_n = \sum_{i=0}^n y_i$ , as  $n \rightarrow \infty$ , then  $S_n \rightarrow y(x)$  so,

$$\|y(x) - S_m\| \leq \frac{\varrho^m}{1-\varrho} \max_{x \in I_0} |y_1|.$$

Thus, in the interval  $I_0$ , the maximum absolute truncation error is

$$\max_{x \in I_0} |y(x) - \sum_{i=0}^m y_i(x)| \leq \frac{\varrho^m}{1-\varrho} \max_{x \in I_0} |y_1|.$$

And thus the proof is finished.

Before starting the application, we will give the following remarks.

**Remark 1** Although the current analysis is conducted in the Banach space  $C([0, T])$ , it is worth noting that the approach may be extended to other functional settings. In particular, weighted spaces and Sobolev spaces  $W^{k,p}$  may provide a more natural framework for problems involving boundary layers or beam-type structures. These spaces allow for more accurate modeling of the regularity and boundary behavior of solutions and could enhance both the theoretical and numerical aspects of the method.

**Remark 2** Equation (15), and equation (16) give many differential operators, for instant, if the differential equation of third order we have six operators by giving the different value of  $n$  and  $m$ .

$$\begin{aligned} L_1(y) &= e^{-i\sqrt{\nu}x} \frac{d}{dx} e^{2i\sqrt{\nu}x} \frac{d}{dx} e^{-i\sqrt{\nu}x} \frac{d}{dx} (y), \\ L_2(y) &= e^{i\sqrt{\nu}x} \frac{d}{dx} e^{-2i\sqrt{\nu}x} \frac{d}{dx} e^{i\sqrt{\nu}x} \frac{d}{dx} (y), \\ L_3(y) &= \frac{d}{dx} e^{-i\sqrt{\nu}x} \frac{d}{dx} e^{2i\sqrt{\nu}x} \frac{d}{dx} e^{-i\sqrt{\nu}x} (y), \\ L_4(y) &= \frac{d}{dx} e^{i\sqrt{\nu}x} \frac{d}{dx} e^{-2i\sqrt{\nu}x} \frac{d}{dx} e^{i\sqrt{\nu}x} (y), \\ L_5(y) &= e^{-i\sqrt{\nu}x} \frac{d}{dx} e^{i\sqrt{\nu}x} \frac{d}{dx} e^{i\sqrt{\nu}x} \frac{d}{dx} e^{-i\sqrt{\nu}x} (y), \\ L_6(y) &= e^{i\sqrt{\nu}x} \frac{d}{dx} e^{-i\sqrt{\nu}x} \frac{d}{dx} e^{-i\sqrt{\nu}x} \frac{d}{dx} e^{i\sqrt{\nu}x} (y). \end{aligned}$$

All of them give the differential equation  $Ly = y''' + \nu y'$ .

**Remark 3** All above differential operators give the same result.

**Proof** We aim to show that all the defined differential operators yield the same result.

Starting with the operator  $L_1(y)$ , we compute:

$$\begin{aligned} L_1(y) &= e^{-i\sqrt{\nu}x} \frac{d}{dx} \left( e^{2i\sqrt{\nu}x} \frac{d}{dx} \left( e^{-i\sqrt{\nu}x} \frac{d}{dx} (y) \right) \right) \\ &= e^{-i\sqrt{\nu}x} \frac{d}{dx} \left( e^{2i\sqrt{\nu}x} \frac{d}{dx} \left( -i\sqrt{\nu} e^{-i\sqrt{\nu}x} y' + e^{-i\sqrt{\nu}x} y'' \right) \right) \\ &= e^{-i\sqrt{\nu}x} \frac{d}{dx} \left( -i\sqrt{\nu} e^{i\sqrt{\nu}x} y' + e^{i\sqrt{\nu}x} y'' \right) \\ &= e^{-i\sqrt{\nu}x} \left( \nu e^{i\sqrt{\nu}x} y' - i\sqrt{\nu} e^{i\sqrt{\nu}x} y'' + i\sqrt{\nu} e^{i\sqrt{\nu}x} y'' \right) \\ &\quad + e^{-i\sqrt{\nu}x} y''' \\ &= e^{-i\sqrt{\nu}x} \left( \nu e^{i\sqrt{\nu}x} y' + e^{-i\sqrt{\nu}x} y''' \right) = y''' + \nu y' \end{aligned}$$

Similarly, it can be shown that:

$$L_2(y) = L_3(y) = L_4(y) = L_5(y) = L_6(y) = y''' + \nu y'$$

Now consider the inverse operator of  $L_1$ :

$$L_1^{-1}(y) = \int_0^x e^{i\sqrt{\nu}x} \left( \int_0^x e^{-2i\sqrt{\nu}x} \left( \int_0^x e^{-i\sqrt{\nu}x} y(x) dx \right) dx \right) dx$$

Applying this operator to  $y''' + \nu y'$ , and using *Mathematica*, we obtain:

$$L_1^{-1}(y'''' + \nu y') = -f(0) + f(x) - \frac{\sin(x\sqrt{\nu})}{\sqrt{\nu}} f'(0) + \frac{-1 + \cos(x\sqrt{\nu})}{\nu} f''(0)$$

Following the same procedure for the inverse operators  $L_2^{-1}, L_3^{-1}, \dots, L_6^{-1}$ , we find that they all yield the same expression:

$$-f(0) + f(x) - \frac{\sin(x\sqrt{\nu})}{\sqrt{\nu}} f'(0) + \frac{-1 + \cos(x\sqrt{\nu})}{\nu} f''(0)$$

Thus, all the differential operators  $L_1, L_2, \dots, L_6$  produce the same result.

Hence, the proof is complete.

## Application

**Example 1** Equation (6) [38–41] with boundary conditions (7) can be written as

$$Ly = -2\alpha \text{Ref}(x) f'(x), \quad (23)$$

we can get  $Ly$  by putting  $m = 0, n = 1$  or  $m = 1, n = 0$  in equation (15),  $m = 0, n = 1$  in equation (16), and  $\nu = (4 - Ha)\alpha^2$ . In this example we used when  $m = 0, n = 1$  in equation (15) that is

$$Ly = e^{\pm i\sqrt{(4-Ha)\alpha^2}x} \frac{d}{dx} e^{\mp 2i\sqrt{(4-Ha)\alpha^2}x} \frac{d}{dx} e^{\pm i\sqrt{(4-Ha)\alpha^2}x} \frac{d}{dx} (y), \quad (24)$$

$L^{-1}$  of equation (24) is

$$\begin{aligned} L^{-1}(y) &= \int_0^x e^{\mp i\sqrt{(4-Ha)\alpha^2}x} \int_0^x e^{\pm 2i\sqrt{(4-Ha)\alpha^2}x} \\ &\quad \int_0^x e^{\mp i\sqrt{(4-Ha)\alpha^2}x} (y) dx dx dx, \end{aligned} \quad (25)$$

applying inverse equation (25) on equation (23), and using Adomian polynomial in equation (21) of non-linear part  $-2\alpha \text{Ref}(x) f'(x)$  we get

$$\begin{aligned} f_0(x) &= 1 + \frac{c}{2}x^2 - \frac{c\alpha^2}{6}x^4 + \frac{cHa\alpha^2}{24}x^4 + \frac{c\alpha^4}{45}x^6 - \frac{cHa\alpha^4}{90}x^6 \\ &\quad + \frac{cHa^2\alpha^4}{720}x^6 - \frac{c\alpha^6}{630}x^8 + \frac{cHa\alpha^6}{840}x^8 - \frac{cHa^2\alpha^6}{3360}x^8 + \frac{cHa^3\alpha^6}{40320}x^8, \\ f_1(x) &= \frac{-c \text{Re}\alpha}{12}x^4 - \frac{c^2 \text{Re}\alpha}{120}x^6 + \frac{c \text{Re}\alpha^3}{45}x^6 - \frac{cHa \text{Re}\alpha^3}{180}x^6 \\ &\quad + \frac{c^2 \text{Re}\alpha^3}{280}x^8 - \frac{c^2 Ha \text{Re}\alpha^3}{1120}x^8 - \frac{c \text{Re}\alpha^5}{420}x^8 \\ &\quad + \frac{cHa \text{Re}\alpha^5}{840}x^8 - \frac{cHa^2 \text{Re}\alpha^5}{6720}x^8, \\ f_2(x) &= \frac{c \text{Re}^2\alpha^2}{180}x^6 + \frac{c^2 \text{Re}^2\alpha^2}{560}x^8 - \frac{c \text{Re}^2\alpha^4}{840}x^8 \\ &\quad + \frac{cHa \text{Re}^2\alpha^4}{3360}x^8, \\ f_3(x) &= -\frac{c \text{Re}^3\alpha^3}{5040}x^8 - \frac{c^2 \text{Re}^3\alpha^3}{5600}x^{10} + \frac{c \text{Re}^3\alpha^5}{28350}x^{10} \\ &\quad - \frac{cHa \text{Re}^3\alpha^5}{113400}x^{10}, \end{aligned}$$

the series solution is given by

$$\begin{aligned}
f(x) = f_0(x) + f_1(x) + f_2(x) + f_3(x) = \\
1 + \frac{cx^2}{2} - \frac{1}{12}cRe\alpha x^4 - \frac{1}{120}c^2Re\alpha x^6 - \frac{1}{6}c\alpha^2 x^4 + \frac{1}{24}cHa\alpha^2 x^4 \\
+ \frac{1}{180}cRe^2\alpha^2 x^6 + \frac{1}{560}c^2Re^2\alpha^2 x^8 + \frac{1}{45}cRe\alpha^3 x^6 - \frac{1}{180}cHaRe\alpha^3 x^6 \\
+ \frac{1}{280}c^2Re\alpha^3 x^8 - \frac{c^2HaRe\alpha^3}{1120}x^8 - \frac{cRe^3\alpha^3}{5040}x^8 - \\
\frac{c^2Re^3\alpha^3}{5600}x^{10} + \frac{1}{45}c\alpha^4 x^6 - \frac{1}{90}cHa\alpha^4 x^6 + \frac{1}{720}cHa^2\alpha^4 x^6 \\
- \frac{1}{840}cRe^2\alpha^4 x^8 + \frac{cHaRe^2\alpha^4}{3360}x^8 - \\
\frac{1}{420}cRe\alpha^5 x^8 + \frac{1}{840}cHaRe\alpha^5 x^8 - \frac{cHa^2Re\alpha^5}{6720}x^8 + \frac{cRe^3\alpha^5}{28350}x^{10} \\
- \frac{cHaRe^3\alpha^5}{113400}x^{10} - \\
\frac{1}{630}c\alpha^6 x^8 + \frac{1}{840}cHa\alpha^6 x^8 - \frac{cHa^2\alpha^6}{3360}x^8 + \frac{cHa^3\alpha^6}{40320}x^8.
\end{aligned}$$

And the solution by SADM as follows

$$\begin{aligned}
f_0(x) &= 1 + \frac{c}{2}x^2, \\
f_1(x) &= -\frac{1}{12}cRe\alpha x^4 - \frac{1}{6}c\alpha^2 x^4 + \frac{1}{24}cHa\alpha^2 x^4 - \frac{1}{120}c^2Re\alpha x^6, \\
f_2(x) &= \frac{1}{180}cRe^2\alpha^2 x^6 + \frac{1}{45}cRe\alpha^3 x^6 - \frac{1}{180}cHaRe\alpha^3 x^6 + \frac{1}{45}c\alpha^4 x^6 \\
&\quad - \frac{1}{90}cHa\alpha^4 x^6 + \frac{1}{720}cHa^2\alpha^4 x^6 + \\
&\quad \frac{1}{280}c^2Re^2\alpha^2 x^8 + \frac{1}{140}c^2Re\alpha^3 x^8 - \frac{1}{560}c^2HaRe\alpha^3 x^8 \\
&\quad + \frac{13}{75600}c^3Re^2\alpha^2 x^{10}, \\
f_3(x) &= \frac{1}{180}cRe^2\alpha^2 x^6 + \frac{1}{90}cRe\alpha^3 x^6 \\
&\quad - \frac{1}{360}cHaRe\alpha^3 x^6 + \frac{13}{10080}c^2Re^2\alpha^2 x^8 + \frac{1}{504}c^2Re\alpha^3 x^8 - \\
&\quad \frac{1}{2016}c^2HaRe\alpha^3 x^8 - \frac{1}{2520}cRe^2\alpha^4 x^8 + \frac{1}{10080}cHaRe^2\alpha^4 x^8 \\
&\quad - \frac{1}{630}cRe\alpha^5 x^8 + \frac{1}{1260}cHaRe\alpha^5 x^8 - \\
&\quad \frac{1}{10080}cHa^2Re\alpha^5 x^8 - \frac{1}{630}c\alpha^6 x^8 + \frac{1}{840}cHa\alpha^6 x^8 \\
&\quad - \frac{1}{3360}cHa^2\alpha^6 x^8 + \frac{1}{40320}cHa^3\alpha^6 x^8 + \\
&\quad \frac{1}{14400}c^3Re^2\alpha^2 x^{10} - \frac{1}{10800}c^2Re^3\alpha^3 x^{10} - \frac{1}{1890}c^2Re^2\alpha^4 x^{10} \\
&\quad + \frac{1}{7560}c^2HaRe^2\alpha^4 x^{10} - \frac{13}{18900}c^2Re\alpha^5 x^{10} + \\
&\quad \frac{13}{37800}c^2HaRe\alpha^5 x^{10} - \frac{13}{302400}c^2Ha^2Re\alpha^5 x^{10} - \\
&\quad \frac{53}{3326400}c^3Re^3\alpha^3 x^{12} - \frac{37}{997920}c^3Re^2\alpha^4 x^{12} \\
&\quad + \frac{37}{3991680}c^3HaRe^2\alpha^4 x^{12} - \frac{89}{165110400}c^4Re^3\alpha^3 x^{14},
\end{aligned}$$

the series solution is given by


$$\begin{aligned}
f(x) = f_0(x) + f_1(x) + f_2(x) + f_3(x) = \\
1 + \frac{c}{2}x^2 - \frac{1}{12}cRe\alpha x^4 - \frac{1}{6}c\alpha^2 x^4 + \frac{1}{24}cHa\alpha^2 x^4 - \\
\frac{1}{120}c^2Re\alpha x^6 + \frac{1}{90}cRe^2\alpha^2 x^6 + \frac{1}{30}cRe\alpha^3 x^6 \\
- \frac{1}{120}cHaRe\alpha^3 x^6 + \frac{1}{45}c\alpha^4 x^6 - \frac{1}{90}cHa\alpha^4 x^6 + \frac{1}{720}cHa^2\alpha^4 x^6 \\
+ \frac{7}{1440}c^2Re^2\alpha^2 x^8 + \frac{23}{2520}c^2Re\alpha^3 x^8 -
\end{aligned}$$

$$\begin{aligned}
&\frac{23}{10080}c^2HaRe\alpha^3 x^8 - \frac{1}{2520}cRe^2\alpha^4 x^8 + \frac{1}{10080}cHaRe^2\alpha^4 x^8 \\
&- \frac{1}{630}cRe\alpha^5 x^8 + \frac{1}{1260}cHaRe\alpha^5 x^8 \\
&- \frac{1}{10080}cHa^2Re\alpha^5 x^8 - \frac{1}{630}c\alpha^6 x^8 + \frac{1}{840}cHa\alpha^6 x^8 - \frac{1}{3360}cHa^2\alpha^6 x^8 \\
&+ \frac{1}{40320}cHa^3\alpha^6 x^8 + \frac{73}{302400}c^3Re^2\alpha^2 x^{10} - \frac{1}{10800}c^2Re^3\alpha^3 x^{10} \\
&- \frac{1}{1890}c^2Re^2\alpha^4 x^{10} + \frac{1}{7560}c^2HaRe^2\alpha^4 x^{10} - \\
&\frac{13}{18900}c^2Re\alpha^5 x^{10} + \frac{13}{37800}c^2HaRe\alpha^5 x^{10} - \frac{13}{302400}c^2Ha^2Re\alpha^5 x^{10} \\
&- \frac{53}{3326400}c^3Re^3\alpha^3 x^{12} - \\
&\frac{37}{997920}c^3Re^2\alpha^4 x^{12} + \frac{37}{3991680}c^3HaRe^2\alpha^4 x^{12} \\
&- \frac{89}{165110400}c^4Re^3\alpha^3 x^{14}.
\end{aligned}$$


Based on the analysis of Figures 3 (a) and (b), which explore the variation of the function ( $f$ ) for ( $\alpha < 0$ ) under different Hartmann numbers ( $Ha$ ) with a fixed Reynolds number ( $Re = 50$ ), the results obtained through NMADM were compared to those derived from the SADM. The comparison indicates a strong agreement between the two methods, with the channel exhibiting a convergent steepness.

In Figures (c) and (d), the focus shifts to the variation of the function ( $f$ ) for ( $\alpha > 0$ ) with different Hartmann numbers while maintaining a fixed Reynolds number ( $Re = 10$ ). Here, a notable difference between the NMADM and SADM results emerges, highlighting a divergent steepness of the channel.


Additionally, Figures 3 (e) and (f) address the variation of the function ( $f$ ) for ( $\alpha > 0$ ) across different Reynolds numbers while keeping the Hartmann number fixed at  $Ha = 5$ . The results indicate a divergence between the NMADM and SADM outcomes, with the steepness of the channel demonstrating divergent behavior in the SADM context, while it remains convergent in the case of NMADM.



(a) SADM at  $Re = 50, \alpha = -5^\circ$  (b) NMADM at  $Re = 50, \alpha = -5^\circ$



(c) SADM at  $Re = 10, \alpha = 7.5^\circ$  (d) NMADM at  $Re = 10, \alpha = 7.5^\circ$



(e) SADM at  $Ha = 50, \alpha = 5^\circ$  (f) NMADM at  $Ha = 50, \alpha = 5^\circ$

Figure 3: NMADM and SADM Solutions for different Hartmann number ( $Ha = 0, 50, 500, 1000, 2000$ ), Reynolds number ( $Re = 10, 15, 25, 35, 50$ ) and ( $\alpha = -5^\circ, 5^\circ, 7.5^\circ$ ).

**Example 2** Consider equation (13) [10–12] with  $EL = 1, p = -2$ , and  $q(x) = -8e^x$ , that is  $\nu = \frac{p}{EL} = -2$  for

$$y^{(4)} - 2y^{(2)} + y = -8e^x, x \in [0, 1], \quad (26)$$

with boundary conditions

$$y(0) = y(1) = 0, y''(0) = 0, y''(1) = -4e. \quad (27)$$

With Exact solution  $y(x) = x(1-x)e^x$ . Equation (26) in an operator form is

$$Ly = -8e^x - y, \quad (28)$$

where  $Ly$  gets when  $m = 0, n = 2$  in equation (16)

$$L(y) = e^{\pm i\sqrt{-2}x} \frac{d}{dx} e^{\mp i\sqrt{-2}x} \frac{d^2}{dx^2} e^{\pm i\sqrt{-2}x} \frac{d}{dx} e^{\pm i\sqrt{-2}x} (y), \quad (29)$$

$L^{-1}$  of equation (29) is

$$L^{-1}(y) = e^{\mp i\sqrt{-2}x} \int_0^x e^{\pm i\sqrt{-2}x} \int_0^x \int_0^x e^{\pm i\sqrt{-2}x} \int_0^x e^{\mp i\sqrt{-2}x} (y) dx dx dx dx. \quad (30)$$

Applying equation (30) to (28) equation we have

$$y_0 = ax + \frac{bx^3}{6} - \frac{x^4}{3} + \left(-\frac{1}{15} + \frac{b}{60}\right)x^5 - \frac{x^6}{30} + \left(-\frac{1}{210} + \frac{b}{1260}\right)x^7 - \frac{x^8}{720} + \left(-\frac{1}{6480} + \frac{b}{45360}\right)x^9 - \frac{x^{10}}{30240},$$

$$y_1 = \frac{-a}{120}x^5 + \left(\frac{-a}{2520} - \frac{b}{5040}\right)x^7 + \frac{x^8}{5040} + \left(\frac{1}{45360} - \frac{a}{90720} - \frac{b}{90720}\right)x^9 + \frac{x^{10}}{90720},$$

$$y_2 = \frac{a}{362880}x^9,$$

$$y_0 + y_1 + y_2 =$$

$$ax + \frac{bx^3}{6} - \frac{x^4}{3} - \frac{ax^5}{120} + \left(-\frac{1}{15} + \frac{b}{60}\right)x^5 - \frac{x^6}{30} + \left(\frac{-a}{2520} - \frac{b}{5040}\right)x^7 + \left(-\frac{1}{210} + \frac{b}{1260}\right)x^7 - \frac{x^8}{840} + \frac{ax^9}{362880} + \left(\frac{1}{45360} - \frac{a}{90720} - \frac{b}{90720}\right)x^9 + \left(-\frac{1}{6480} + \frac{b}{45360}\right)x^9 - \frac{x^{10}}{45360}.$$

Using boundary conditions in equation (27) in the above series solution we get  $a = 1$ , and  $b = -3$ .

$$y_0 + y_1 + y_2 = x - \frac{x^3}{2} - \frac{x^4}{3} - \frac{x^5}{8} - \frac{x^6}{30} - \frac{x^7}{144} - \frac{x^8}{840} - \frac{x^9}{5760} - \frac{x^{10}}{45360}.$$

The Solution by SADM

$$y_0 = ax + \frac{bx^3}{6} - \frac{x^4}{3} - \frac{x^5}{15} - \frac{x^6}{90} - \frac{x^7}{630} - \frac{x^8}{5040} - \frac{x^9}{45360} - \frac{x^{10}}{453600},$$

$$y_1 = \left(\frac{-a}{120} + \frac{b}{60}\right)x^5 - \frac{x^6}{45} + \left(-\frac{1}{315} - \frac{b}{5040}\right)x^7 - \frac{x^8}{5040} - \frac{x^9}{45360} - \frac{x^{10}}{453600},$$

$$y_2 = \left(\frac{-a}{2520} + \frac{b}{1260}\right)x^7 - \frac{x^8}{1260} + \left(-\frac{1}{11340} + \frac{a}{362880} - \frac{b}{90720}\right)x^9,$$

$$y_0 + y_1 + y_2 = ax + \frac{bx^3}{6} - \frac{x^4}{3} - \frac{x^5}{15} + \left(\frac{-a}{120} + \frac{b}{60}\right)x^5 - \frac{x^6}{30} - \frac{x^7}{630} + \left(-\frac{1}{315} - \frac{b}{5040}\right)x^7 + \left(\frac{-a}{2520} + \frac{b}{1260}\right)x^7 - \frac{x^8}{840} - \frac{x^9}{22680} + \left(-\frac{1}{11340} + \frac{a}{362880} - \frac{b}{90720}\right)x^9 - \frac{x^{10}}{226800}.$$

Using boundary conditions in equation (27) in the above series solution we get  $a = 1$ , and  $b = -3$ .

$$y_0 + y_1 + y_2 = x - \frac{x^3}{2} - \frac{x^4}{3} - \frac{x^5}{8} - \frac{x^6}{30} - \frac{x^7}{144} - \frac{x^8}{840} - \frac{x^9}{5760} - \frac{x^{10}}{45360}.$$

The following is the Taylor series for the exact solution

$$x - \frac{x^3}{2} - \frac{x^4}{3} - \frac{x^5}{8} - \frac{x^6}{30} - \frac{x^7}{144} - \frac{x^8}{840} - \frac{x^9}{5760} - \frac{x^{10}}{45360} + O(x^{11}).$$

To clarify the convergence between the exact solution and the solutions obtained using the SADM and the NMADM in this example, we utilize a tenth-order Taylor series expansion. We find that the solution provided by the NMADM is identical to the exact solution and is closer to the exact solution than that of the SADM.

**Example 3** Consider a special sixth-order boundary value problem with parameter  $c$  [42–44]

$$y^{(6)}(x) - (1+c)y^{(4)}(x) = cx - cy^{(2)}(x), \quad (31)$$

with boundary conditions

$$y(0) = y'(0) = 1, y''(0) = 0,$$

$$y(1) = \frac{7}{6} + \sinh(1), y'(1) = \frac{1}{2} \cosh(1), y''(1) = 1 + \sinh(1). \quad (32)$$

$y(x) = 1 + \frac{1}{6}x^3 + \sinh(x)$  is the exact solution. Equation (31) in an operator form is

$$Ly = cx - cy^{(2)}(x), \quad (33)$$

$L$  obtains when  $m = 3, n = 1$  and  $\nu = -(1+c)$  in equation (15)

$$Ly = \frac{d^3}{dx^3} e^{\pm i \sqrt{-(1+c)} x} \frac{d}{dx} e^{\mp 2i \sqrt{-(1+c)} x} \frac{d}{dx} e^{\pm i \sqrt{-(1+c)} x} \frac{d}{dx} (y), \quad (34)$$

and  $L^{-1}$  of equation (34)

$$L^{-1} = \int_0^x e^{\mp i \sqrt{-(1+c)} x} \int_0^x e^{\pm 2i \sqrt{-(1+c)} x} \int_0^x e^{\mp i \sqrt{-(1+c)} x} dx dx dx dx dx dx (y). \quad (35)$$

Applying (35) on (33) we get

$$\begin{aligned} y_0 &= 1 + x + \frac{x^3 a_1}{6} + \frac{x^4 a_2}{24} + \frac{x^5 a_3}{120} + \frac{x^6 a_2}{720} + \frac{cx^6 a_2}{720} + \frac{cx^7}{5040} \\ &\quad + \frac{x^7 a_3}{5040} + \frac{cx^7 a_3}{5040} + \frac{x^8 a_2}{40320} + \frac{cx^8 a_2}{20160} + \frac{c^2 x^8 a_2}{40320} + \\ &\quad \frac{cx^9}{362880} + \frac{x^9 a_3}{362880} + \frac{cx^9 a_3}{181440} + \frac{c^2 x^9 a_3}{362880} + \frac{c^2 x^9}{362880} + \frac{x^{10} a_2}{3628800} + \\ &\quad \frac{cx^{10} a_2}{1209600} + \frac{c^2 x^{10} a_2}{1209600} + \frac{c^3 x^{10} a_2}{3628800}, \\ y_1 &= -\frac{cx^7 a_1}{5040} - \frac{cx^8 a_2}{40320} - \frac{cx^9 a_1}{362880} - \frac{c^2 x^9 a_1}{362880} - \frac{cx^9 a_3}{362880} \\ &\quad - \frac{cx^{10} a_2}{1814400} - \frac{c^2 x^{10} a_2}{1814400}, \end{aligned}$$

the series solution by NMADM

$$\begin{aligned} y(x) &= y_0 + y_1 \\ &= 1 + x + \frac{x^3 a_1}{6} + \frac{x^4 a_2}{24} + \frac{x^5 a_3}{120} + \frac{x^6 a_2}{720} + \frac{cx^6 a_2}{720} + \frac{cx^7}{5040} \\ &\quad - \frac{cx^7 a_1}{5040} + \frac{x^7 a_3}{5040} + \frac{cx^7 a_3}{5040} + \frac{x^8 a_2}{40320} + \frac{cx^8 a_2}{40320} \\ &\quad + \frac{c^2 x^8 a_2}{40320} + \frac{cx^9}{362880} + \frac{cx^9 a_3}{362880} + \frac{c^2 x^9 a_3}{362880} - \frac{cx^9 a_1}{362880} - \frac{c^2 x^9 a_1}{362880} + \frac{x^9 a_3}{362880} \\ &\quad + \frac{cx^9 a_3}{362880} + \frac{c^2 x^9 a_3}{362880} + \frac{x^{10} a_2}{3628800} + \frac{cx^{10} a_2}{3628800} + \frac{c^2 x^{10} a_2}{3628800} + \frac{c^3 x^{10} a_2}{3628800}. \end{aligned}$$

And the solution in SADM

$$\begin{aligned} y_0 &= 1 + x + \frac{x^3 a_1}{6} + \frac{x^4 a_2}{24} + \frac{x^5 a_3}{120} + \frac{cx^7}{5040}, \\ y_1 &= \frac{x^6 a_2}{720} + \frac{cx^6 a_2}{720} - \frac{cx^7 a_1}{5040} + \frac{x^7 a_3}{5040} + \frac{cx^7 a_3}{5040} + \frac{cx^9}{362880} \\ &\quad + \frac{c^2 x^9}{362880} - \frac{cx^8 a_2}{40320} - \frac{cx^9 a_3}{362880} - \frac{c^2 x^{11}}{39916800}, \end{aligned}$$

$$\begin{aligned} y_0 + y_1 &= 1 + x + \frac{x^3 a_1}{6} + \frac{x^4 a_2}{24} + \frac{x^5 a_3}{120} + \frac{x^6 a_2}{720} \\ &\quad + \frac{cx^6 a_2}{720} + \frac{cx^7}{5040} - \frac{cx^7 a_1}{5040} + \frac{cx^7 a_3}{5040} \\ &\quad - \frac{cx^8 a_2}{40320} + \frac{cx^9}{362880} + \frac{c^2 x^9}{362880} - \frac{cx^9 a_3}{362880} - \frac{c^2 x^{11}}{39916800}. \end{aligned}$$

The value of constants  $a_1, a_2$  and  $a_3$  can be get by using boundary conditions in (32).




Figure 4: Comparing between Exact, SADM and NMADM solution when  $y(x) = y_0 + y_1$  for different value of  $c = 0, 20, 100, 1000$

The performance of the new modified Adomian decomposition method (NMADM) and the Standard Adomian Decomposition Method (SADM) reveals significant differences as the parameter ( $c$ ) increases. From our analysis, particularly illustrated in Figure 4, it is evident that NMADM consistently provides solutions that closely approximate the exact solution, irrespective of how large the parameter ( $c$ ) becomes. This robustness indicates the effectiveness of NMADM in handling higher values of ( $c$ ) without significant deviation from the true solution. In stark contrast, the SADM exhibits a noticeable divergence from the exact solution as the parameter ( $c$ ) increases. This indicates that while SADM may perform adequately for lower values of ( $c$ ), its reliability diminishes significantly as the parameter grows. Furthermore, our findings align with previous literature [43], which has stated that the approximate solutions derived from the Adomian Decomposition Method are quite accurate for small values of ( $c$ ) only. As ( $c$ ) increases, the accuracy of the SADM solutions deteriorates, highlighting a limitation in this method, which NMADM effectively overcomes.

**Remark 4** While the focus of this study has been on higher-order ordinary differential equations, the structure of the modified operator introduced in the NMADM suggests potential applicability to certain classes of partial differential equations (PDEs). In particular, extension to nonlinear diffusion–reaction PDEs may be feasible, given the operator's capability to handle nonlinearity and higher-order derivatives. This opens a possible direction for future investigation.



## Conclusion

The study presented a new modified Adomian decomposition method (NMADM) tailored for solving higher-order ordinary differential equations, particularly in the context of magnetohydrodynamic (MHD) flows, elastic beam dynamics, and sixth-order boundary value problems in physics. The key findings of this research indicate that the modified approach demonstrates significant advantages over traditional methods in terms of convergence speed and accuracy. By applying the NMADM, the researchers were able to achieve rapid convergence for various complex problems, providing explicit analytical solutions that can effectively describe physical phenomena in MHD flows and the behavior of elastic beams. The method's versatility also extends to accommodating sixth-order boundary value problems, which are often encountered in engineering and applied physics.

In addition to the analytical benefits, the structure of NMADM suggests favorable scalability to more complex or higher-dimensional problems. The incorporation of weighted exponential operators enhances the convergence behavior, which may reduce computational costs in multidimensional applications. Furthermore, its semi-analytical nature avoids the intensive discretization required by many numerical methods. Future studies could explore its integration with domain decomposition or spectral methods to improve performance in large-scale simulations. The results validate the efficiency of the modified Adomian decomposition method as a robust computational tool, suggesting its potential for broader applications in tackling nonlinear dynamic systems and enhancing the understanding of diverse physical mechanisms. Further studies could explore its applicability to other types of differential equations and complex physical models, paving the way for advancements in both theoretical and practical realms of physics.

## Ethics approval and consent to participate

Not applicable

## Consent for publication

Not applicable

## Availability of data and materials

The raw data required to reproduce these findings are available in the body and illustrations of this manuscript.

## Author's contribution

The authors confirm contribution to the paper as follows: Study conception and design: Nuha Mohammed Dabwan; Theoretical calculations and modeling: Nuha Mohammed Dabwan; Data analysis and validation: Nuha Mohammed Dabwan, Yahya Qaid Hasan; Draft manuscript preparation: Nuha Mohammed Dabwan. Both authors reviewed the results and approved the final version of the manuscript.

## Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

## Conflicts of interest

The authors declare that there is no conflict of interest regarding the publication of this article.

## Acknowledgements

The authors would like to express their sincere gratitude to the anonymous reviewers for their valuable comments and constructive suggestions, which have significantly improved the quality of this manuscript. The authors also extend their appreciation to the editorial board of An-Najah University Journal for Research – A for their professional handling and continuous support during the peer-review process.

## Open Access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <https://creativecommons.org/licenses/by-nc/4.0/>

## References

- [1] Mikusinski J. *The Operational Calculus*. Oxford: Pergamon Press; 1959.
- [2] Oldham KB, Spanier J. *The Fractional Calculus: Theory and Applications of Differentiation and Integration to Arbitrary Order*. Academic Press; 1974.
- [3] Samko SG, Kilbas AA, Marichev OI. *Fractional Integrals and Derivatives: Theory and Applications*. Gordon and Breach Science Publishers; 1993.
- [4] Jeffery GB. The two-dimensional steady motion of a viscous fluid. *Philos Mag*. 1915;6:455–465.
- [5] Hamel G. Spiralförmige Bewegungen Zäher Flüssigkeiten. *Jahresber Dtsch Math-Ver*. 1916;25:34–60.
- [6] Bansal L. *Magnetofluidynamics of Viscous Fluids*. Jaipur: Jaipur Publishing House; 1994.
- [7] Esmaili Q, Ramiar A, Alizadeh E, Ganji DD. An approximation of the analytical solution of the Jeffery-Hamel flow by decomposition method. *Phys Lett A*. 2008;372:3434–3439.
- [8] Elishakoff I. Essay on the contributors to the elastic stability theory. *Meccanica*. 2005;40:75–110. doi:10.1007/s11012-004-2199-y
- [9] Chen H, Cui Y. Existence and uniqueness of solutions to the nonlinear boundary value problem for fourth-order differential equations with all derivatives. *J Inequal Appl*. 2023;2023:23. doi:10.1186/s13660-022-02907-9.
- [10] Huang W, Jiang T, Zhang X, Khan NA, Sulaima M. Analysis of beam-column designs using a new soft computing technique. *Complexity*. 2021;Article ID 6639032. doi:10.1155/2021/6639032.
- [11] Kelesoglu O. The solution of fourth order boundary value problem using Adomian decomposition method. *Math Probl Eng*. 2014;Article ID 649471. doi:10.1155/2014/649471.
- [12] Taiwo OA, Ogunlaran OM. A non-polynomial spline method for solving linear fourth-order boundary-value problems. *Int J Phys Sci*. 2011;6(13):3246–3254.
- [13] Khalid A, Naeem MN, Agarwal P, Ghaffar A, Ullah Z, Jain S. Numerical approximation for the solution of linear sixth order boundary value problems by cubic B-spline. *Adv Differ Equ*. 2019;2019:492.

- [14] Khalid M, Sultana M, Zaidi F. Numerical solution of sixth order differential equations arising in astrophysics by neural network. *Int J Comput Appl*. 2014;107(6).
- [15] Adomian G. *Solving Frontier Problem of Physics: The Decomposition Method*. London: Kluwer Academic Publishers; 1994.
- [16] Adomian G. A review of the decomposition method in applied mathematics. *J Math Anal Appl*. 1988;135:501–544.
- [17] Adomian G, Rach R. Modified Adomian polynomials. *Math Comput Model*. 1996;24(11):39–46.
- [18] Alomari SN, Hasan YQ. Solving the hyperbolic telegraph equation using a modified Adomian decomposition method. *Asian Res J Math*. 2023;19(9):278–293.
- [19] Dabwan NM, Hasan YQ. Adomian method for solving Emden-Fowler equation of higher order. *Adv Math Sci J*. 2020;9(1):231–240.
- [20] Dabwan NM, Hasan YQ. An approximate solution of higher order ordinary differential equations using modified Adomian decomposition method. *IJRDO J Math*. 2020;6:1–8.
- [21] Dabwan NM, Hasan YQ. Solving second order ordinary differential equation using a new modified Adomian method. *Adv Math Sci J*. 2020;3(9):937–943.
- [22] Hasan YQ. Modified Adomian decomposition method for solving higher odd order boundary value problems. *MAYFEB J Math*. 2017;2:74–79.
- [23] Hasan YQ, Dabwan NM, Alaqel SA, Othman SG, AL-Rabahi ZA. Using Adomian decomposition method for solving systems of second order ordinary differential equation. *J Math Comput Sci*. 2021;11(2):2204–2217.
- [24] Othman SG, Hasan YQ. Adomian decomposition method for solving oscillatory systems of higher order. *Adv Math Sci J*. 2020;3(9):945–953.
- [25] Othman SG, Hasan YQ. Solving ordinary differential equation of higher order by Adomian decomposition method. *Asian J Probab Stat*. 2020;9(3):44–53.
- [26] Wazwaz AM. A reliable modification of Adomian decomposition method. *Appl Math Comput*. 1999;102:77–86.
- [27] Wazwaz AM. A new algorithm for calculating Adomian polynomials for nonlinear operators. *Appl Math Comput*. 2000;111:53–69.
- [28] Odiba OJ. The modified Adomian decomposition method for the solution of third order ordinary differential equations. *IOSR J Math*. 2017;13(6):61–64.
- [29] Hashim I. Adomian decomposition method for solving BVPs for fourth-order integro-differential equations. *J Comput Appl Math*. 2006;193:658–664.
- [30] Ali N, Ahmad S, Aziz S, Zaman G. The Adomian decomposition method for solving HIV infection model of latently infected cells. *Matrix Sci Math*. 2019;3(1):5–8.
- [31] Coşkun SB, Öztürk B, Mutman U. Adomian decomposition method for vibration of nonuniform Euler beams on elastic foundation. *Proc 9th Int Conf Struct Dyn*. 2014:1935–1940.
- [32] Bougoffa L, Mziou S, Rach RC. Exact and approximate analytical solutions of the Jeffery-Hamel flow problem. *Math Model Anal*. 2016;21(2):174–187.
- [33] Chen WF, Atsuta T. *Theory of Beam-Columns. Vol. 2: Space Behavior and Design*. New York: McGraw-Hill; 1977.
- [34] Timoshenko S, Gere JM. *Theory of Elastic Stability*. Mineola, NY: Dover Publications; 1961.
- [35] Timoshenko SP, Gere JM. *Theory of Elastic Stability*. Chelmsford, MA: Courier Corporation; 2009.
- [36] Duan JS, Rach RR. A new modification of the Adomian decomposition method and its convergence. *Appl Math Comput*. 2012;218(19):10042–10054.
- [37] Wazwaz AM, El-Sayed SM. A new modification of the Adomian decomposition method. *Appl Math Comput*. 2001;122:393–405.
- [38] Abbasbandy S, Shivanian E. Exact analytical solution of the MHD Jeffery-Hamel flow problem. *Meccanica*. 2012;47:1379–1389.
- [39] Domairry DG, Mohsenzadeh A, Famouri M. The application of homotopy analysis method to solve nonlinear differential equation governing Jeffery-Hamel flow. *Commun Nonlinear Sci Numer Simul*. 2009;14:85–95.
- [40] Ganji DD, Sheikholeslami M, Ashorynejad HR. Analytical approximate solution of nonlinear differential equation governing Jeffery-Hamel flow. *ISRN Math Anal*. 2011;Article ID 937830. doi:10.5402/2011/937830
- [41] Sheikholeslami M, Ganji DD, Ashorynejad HR, Rokni HB. Analytical investigation of Jeffery-Hamel flow. *Appl Math Mech*. 2012;33(1):25–36.
- [42] GUL M, KHAN H, ALI A. The solution of fifth and sixth order linear and nonlinear boundary value problems by the improved residual power series method. *J Math Anal Model*. 2022;3(1):1–14.
- [43] Momani S, Noor MA, Mohyud-Din ST. Numerical methods for solving a special sixth-order boundary value problem. *Nonlinear Anal Forum*. 2008;13(2):1–10.
- [44] Noor MA, Noor KI, Mohyud-Din ST. Variational iteration method for solving sixth-order boundary value problems. *Commun Nonlinear Sci Numer Simul*. 2009;14:2571–2580.

Feature	NMADM	Duan-Rach Modification [36]	Wazwaz Modification [37]
Linear Operator (L)	Redefined linear operator $L(y) = y^{(r+2)} + \nu y^{(r)}$ , reformulated using integral-exponential operators for improved spectral condition and stability.	Traditional operator $L(y) = y^{(r+2)}$ with adjustments for boundary conditions.	Second-order operator designed to simplify nonlinear expressions.
Key Innovation	Redesign of the operator structure to enhance spectral properties and convergence in stiff or oscillatory problems.	Improved inversion matching specific boundary conditions.	Utilization of He polynomials for nonlinear term approximation.
Convergence Enhancement	Faster convergence due to better operator conditioning and exponential-integral formulation.	Improved convergence for problems with structured boundaries.	Improved convergence by matching structure of nonlinear terms.
Typical Applications	Stiff and oscillatory nonlinear differential equations.	Nonlinear boundary value problems and weak nonlinear systems.	PDEs and solitary wave solutions.
Distinct Advantage	Improved conditioning and accelerated convergence.	Flexible inversion options.	Compatible with analytical tools like He's method.
Proposed Improvement	Customized operator based on the problem's structure, controlling higher-order term growth.	Focus on operator inversion only, no structural changes to the equation.	Alteration of nonlinear terms' form, base equation remains unchanged.

Table 1: Comparison between NMADM and other Modified ADM Variants

$Ha$ (Hartmann number)	$c = f''(0)$			
	$\alpha = -5^\circ$		$\alpha = 5^\circ$	
	NMADM	SADM	NMADM	SADM
0	-1.78545	-1.783225	-2.25333	-2.25429
200	-1.57552	-1.572131	-1.98479	-1.98461
400	-1.39632	-1.38969	-1.75465	-1.75529
600	-1.23815	-1.23185	-1.55609	-1.55813
800	-1.10261	-1.09500	-1.38393	-1.38726
1000	-0.98481	-0.976044	-1.23409	-1.23828
2000	-0.583545	-187.87	-1.72186	-0.72328

Table 2: **(Example 1)**  $c = f''(0)$  at different  $Ha$ ,  $Re=10$ ,  $\alpha = 5^\circ$  and  $\alpha = -5^\circ$ .

	SADM			NMADM		
	$c=0$	$c=20$	$c=100$	$c=0$	$c=20$	$c=100$
$a_1$	2.00411	2.00293	1.84544	2.00000313	2.0000061	1.99999991
$a_2$	-0.0441241	0.237589	1.01826	-0.00000288	-0.0006482	0.0000002
$a_3$	1.16996	-0.12382	-5.70828	1.00009	1.00033	0.999997

Table 3: **(Example 3)** Comparing value of  $y(1) = a_1$ ,  $y'(1) = a_2$  and  $y''(1) = a_3$  at different  $c$  using SADM and NMADM when  $y(x) = y_0 + y_1$