

**BASE CLEAVAGE OF SOME BENZYL–SILICON AND
TIN COMPOUNDS IN H₂O – MeOH – CYCLODEXTRIN MEDIUM**

BY

FOAD M. S. MAHMOUD

Department of chemistry , An - Najah National University

Nablus , West Bank .

المخلص

لقد تمت دراسة تأثير ألفا - وبيتا - سيكلوديكتريز على معدل سرعة التفاعل بواسطة قاعدة على
3, 5 - Cl₂. C₆ H₃ CH₂. MMe₃ , 4 - ClC₆H₄ CH₂ MMe₃. 3 - ClC₆ H₄ CH₂ MMe₃
حيث ان M = Si , Sn . ألفا - وبيتا - سيكلوديكتريز تزيد من معدل سرعة التفاعل في حالة مركبات
القصدير RSnMe₃ ولكن لا يوجد اي حفز ذو مغزى في حالة مركبات السيليكون المشابهة RSiMe₃

ABSTRACT


The effects of α - and β - cyclodextrin (α - , β - CD's) on the rates of base cleavage of 4 - CNC₆ H₄ SnMe₃ , 3-ClC₆ H₄ CH₂ MMe₃ , 4 - ClC₆ H₄ CH₂- MMe₃ , and 3,5 - Cl₂ C₆ H₃ CH₂ MMe₃ with M = Si and Sn have been examined in H₂O – MeOH medium . α - and β -CD's accelerate the cleavage of the RSnMe₃ compounds (R=3- ClC₆ H₄ CH₂ , 4-ClC₆ H₄ CH₂, and 3,5 - Cl₂ C₆H₃ CH₂) . There is no significant catalysis in the case of the corresponding silicon compounds .

INTRODUCTION


Extensive studies have been made of base cleavage of R-MMe₃ bonds , where M = Si or Sn and R is an organic group , usually of the aryl or benzyl type [1-9]. The medium of NaOMe-MeOH has been most frequently used , and in it the cleavages of the silicon and tin compounds have been shown to differ in mechanism ; for the silicon compounds the rate determining step involves the separation of the carbanion R⁻ mechanism B (or B_s), whereas for most tin compounds the breaking of the Sn - R bond is accompanied by proton transfer (electrophilic assistance) from the solvent to the separating carbon center [5-8] mechanism A (or A_s). The exceptions for tin compounds arise in the case of very reactive species , for example with R = 9-fluorenyl , which appear to undergo cleavage without electrophilic assistance [8] mechanism B (or B_s).

The following mechanisms were suggested [7,11,12,13] for cleavage of some benzyltrimethylsilanes and stannanes by methanolic sodium methoxide :


1) A.




2) A_s.



3) B.



4) B_s.



For cleavage of Benzylsilicon compounds, in a medium of fairly high water content (36 vol %), the value of the specific rate constant, k_s , given by the observed first-order rate constant divided by the base concentration, is fairly constant over a wide range of base concentration. Whereas in methanol depends markedly on the sodium methoxide concentration [10].

EXPERIMENTAL

Preparation of organosilicon and organotin compounds

The $\text{XC}_6\text{H}_4\text{CH}_2\text{SiMe}_3$ compounds were made by standard methods, viz. (X =), 3-Cl [23], 4-Cl [24]. 4-ClC₆H₄CH₂SnMe₃ was made by the standard method²⁴

The following compounds were kindly supplied by Professor C. Eaborn:

(3,5-Cl₂C₆H₃CH₂SiMe₃) 3,5-Cl₂C₆H₃CH₂SnMe₃, 3-ClC₆H₄CH₂SnMe₃
4-ClC₆H₄CH₂SnMe₃.

Rate measurements

The rates of the cleavage of organosilicon, and organotin compounds were measured spectrophotometrically using a Unicam SP 1700 with automatic recording system. The machine was fitted with a thermostatically controlled cell-holder accurate to within $\pm 0.1^\circ$.

Cyclodextrin solutions were prepared as follows. The CD was dissolved in water (5 ml) and (4 ml) sample of the sodium methoxide solution in methanol was then added. A (4 ml) sample of the resulting solution was added to (1 ml) of the solution of the organometallic in methanol. The mixture was shaken and transferred to the stoppered quartz cell (10 mm path length) and the change in UV absorption at the selected wavelength was recorded. The reference cell was filled with aqueous methanol.

Optical density readings were recorded at convenient intervals up to three half-lives. The infinity readings were taken after ten half-lives. The spectra were found to be in good agreement with solutions of authentic samples of the appropriate toluene at the correct concentration in the cleavage medium.

The following wavelengths were used in the rate studies:

3,5 - Cl₂ - C₆H₃ CH₂ SnMe₃, 280 ; 3,5 - Cl₂ C₆H₃ SiMe₃, 283.5 ;

3 - ClC₆H₄CH₂ SnMe₃, 286 ; 3 - ClC₆H₄CH₂ SiMe₃, 279.5 ;

4 - ClC₆H₄CH₂SnMe₃, 286 ; 4 - ClC₆H₄CH₂SiMe₃, 278 ; 4 - CNC₆H₄CH₂ SnMe₃, 270 nm .

RESULTS

We have now examined the cleavages of 3,5 - Cl₂C₆H₃MMe₃, 3 - Cl- C₆H₄CH₂MMe₃, 4 - ClC₆H₄CH₂MMe₃, and 4 - CNC₆H₄CH₂SnMe₃ catalysed by α - and β -CD and the results are shown in Tables 1 and 2 .

Table 1

Rates of cleavage of RMMe_3 by $\text{NaOH-H}_2\text{O-MeOH-}\infty$ - CD at 50°

R	$10^3[\infty\text{-CD}]/\text{M}$	$\text{H}_2\text{O:MeOH}$ by volume	$[\text{NaOH}]^*/\text{M}$	RSnMe_3 10^5ks/l $\text{mol}^{-1}\text{s}^{-1}$	RSiMe_3 10^5ks/l $\text{mol}^{-1}\text{s}^{-1}$
3,5- $\text{Cl}_2\text{C}_6\text{H}_3\text{CH}_2$	0	44:56	0.238	913.33	109.50
	41.79	44:56	0.238	2206.33	127.73
	26.82	44:56	0.031	4386.67	—
	0	44:56	0.031	1155.17	—
3- $\text{ClC}_6\text{H}_4\text{CH}_2$	0	44:56	0.953	26.72	2.72
	41.79	44:56	0.953	75.77	2.35
4- $\text{ClC}_6\text{H}_4\text{CH}_2$	0	44:56	0.953	8.33	0.43
	41.79	44:56	0.953	20.20	0.48

★ The base is denoted throughout by NaOH although both hydroxide and methoxide ions are present .

Table 2

Rates of cleavage of RMe_3 by $\text{NaOH-H}_2\text{O-MeOH-}\beta\text{-CD}$ at 50° .

R	$10^3 \cdot [\beta\text{-CD}]/\text{M}$	$\text{H}_2\text{O:MeOH}$ by volume	$[\text{NaOH}]/\text{M}$	RMe_3 , 10^5ks/l $\text{mol}^{-1}\text{s}^{-1}$.	RSiMe_3 , 10^5ks/l $\text{mol}^{-1}\text{s}^{-1}$.
3,5- $\text{Cl}_2\text{C}_6\text{H}_3\text{CH}_2$.	3.69	23.8:76.2	0.306	1794.0	118.65
	38.50	44:56	0.238	2116.5	84.93
	58.03	44:56	0.257	2197.2	96.67
	22.03	44:56	0.031	5333.3	---
	13.66	44:56	0.052	3219.7	---
	22.03	44:56	0.100	2888.2	---
	15.86	44:56	0.205	2336.5	---
	44.05	44:56	0.482	1997.3	---
	0	44:56	0.031	1155.2	---
	0	44:56	0.100	837.0	---
	0	44:56	0.205	983.8	---
3- $\text{ClC}_6\text{H}_4\text{CH}_2$.	0	44:56	0.953	26.72	2.72
	28.4	37.5:62.5	1.089	61.67	2.98
	39.5	44:56	0.953	80.00	2.07
	44.05	44:56	0.483	59.50	---
4- $\text{CNC}_6\text{H}_4\text{CH}_2$.	0	0:100	0.007 *	66666.7	---
	7.36	44:56	0.007	64116.7	---

* The base is NaOMe.

It is seen from Table 1 that α -CD catalyses the cleavage of 3,5-Cl₂C₆H₃CH₂-SnMe₃, 3-Cl-C₆H₄CH₂SnMe₃, and 4-Cl-C₆H₄CH₂SnMe₃ by a factor of Ca. 2.5 at a moderate base concentration and by the larger factor of Ca. 3.8 at a lower base concentration [0.03 M]. For the corresponding silicon compounds there is a little or no acceleration by α -CD.

In Table 2 results are listed for the addition of β -CD to the cleaving medium, and it will be seen that the effects are very similar.

The lower rates for cleavage of some organosilicon compounds in the presence rather than in the absence of β -CD as shown in Table 2.

For the cleavage of 4-CNC₆H₄CH₂SnMe₃ there is no catalysis by β -CD as shown in Table 2.


There has been no change in the ultraviolet spectrum of 3-Cl-C₆H₄CH₂MMe₃ or 3,5-Cl₂C₆H₃CH₂MMe₃ when α - or β -CD was added.

DISCUSSION

Cyclodextrins are fairly stable in alkaline solution, but are rather susceptible to acid catalysis [14].

Both α - and β -CD's accelerate the cleavage of a variety of phenyl esters. The catalysis by CD's varies widely, from 300 fold for *m*-nitrophenyl acetate to 1.1 fold for *t*-butylphenyl acetate [15].

The α -CD catalysed hydrolysis of *p*-nitrofluoroacetanilide is thought to involve the process shown below [16]:




As shown in Tables 1 and 2 there is a difference in the behaviour of tin and silicon compounds. Catalysis by both α and β -CD's is covalent which was observed in the cleavage of phenyl acetates and acetanilides [15]. In this catalysis the secondary hydroxyl group (in the ionic stage) will provide the nucleophile to attack the metal atom in the compounds shown in Tables 1 and 2. Evidence for the extensive ionization of the secondary hydroxyl groups (pK_a 12) [17-19] is provided by the observation that during the preparation of the cleavage systems, the solubility of both α and β -CD's in aqueous methanol was increased by addition of base (NaOMe).

The higher catalysis by α - and β -CD's catalysed cleavages of the organotin compound (3,5-Cl₂C₆H₃CH₂SnMe₃) at low base concentration is in support of electrophilic assistance by the unionized secondary hydroxyl groups which are more available at low than at higher base concentration.

The lower rates of some organosilicon compounds in the presence rather than in the absence of β -CD are possibly due to the fact that some NaOMe will be used in the ionization of secondary hydroxyl groups of β -CD and the resulting anions will be very poor nucleophiles.


The above results are consistent with the electrophilic assistance by solvent (in CD's it is by the unionized secondary hydroxyl groups) in the rate-determining step in the cleavage of RSnMe₃ compounds



For the cleavage of 4-CN C₆H₄CH₂SnMe₃ there is no catalysis by β -CD as shown in Table 2. This suggests that this compound possibly reacts by the ionization mechanism B (or B_s), that is the mechanism which operates for the cleavage of benzylsilicon compounds, and it is significant that Eaborn et al [8] suggested that the mechanistic crossover point for cleavages of XC₆H₄CH₂SnMe₃ compounds in MeOH-MeONa lies near the compound with X = 4-CN.

The possibility was considered that it might be the SnMe₃ group and not the aryl group which enters the cavity of the CD, but a simple calculation indicated that the diameter of this group (viz. 7.55 Å) was markedly greater than that of the cavity (viz. 7 Å) [20]. However calculation also indicated that the width of the 3,5-Cl₂C₆H₃CH₂ position of the 3,5-Cl₂C₆H₃CH₂MMe₃ (viz. 9.12 Å)

is also too great to permit full entry into the cavity ; however this would also be the case for the 3,5 - Me₂C₆H₃ - group , and yet both α - and β - CD's are shown to catalyse the reactions of dimethylphenyl acetates [21] . It is conceivable that only part of the 3,5 - Cl₂C₆H₃ - group enters the cavity , as shown below .



It is relevant in this connection to note that in hydrolysis of phenyl esters , meta - substituted phenyl esters show larger acceleration by CD than corresponding para - compound , and meta / para specificity is larger for esters with more bulky substituents . These are due to the geometry of the inclusion complexes , especially the distance between the electrophilic centers (the carbon atom in phenyl esters) and the nucleophilic catalytic center (a secondary hydroxyl group of the CD) .^[22]

Finally we note that we obtained no change in the ultraviolet spectrum of 3 - ClC₆H₄CH₂MMe₃ or 3,5 - Cl₂C₆H₃CH₂MMe₃ when α - or β - CD was added . This indicates that there is only a small degree of inclusion even in the case of the tin compounds , which is perhaps not surprising since the organic group will have a substantially greater affinity for these methanolic media than for the aqueous media usually used with CD'S , and so shows a lower tendency to enter the non-polar cavity . Since the degree of inclusion is so small , in our media , this implies that the reactivity of an included molecule must be exceptionally high to give rise to the marked catalysis observed .

ACKNOWLEDGEMENT

We thank Professor C . Eaborn for the gift of some organotin and organosilicon compounds .

REFERENCES

- [1] Eaborn C. and Parker S. , (1955) , J. Chem. Soc. 126 .
- [2] Eaborn C. , Sinnitambe K. and Walton D., (1975) , J. Chem. Soc. Perkin II 380 .
- [3] Bott R, Eaborn C. and Swaddle T., (1972) , J. Chem. Soc. 2342 .
- [4] Eaborn C., Najam A. and Walton D., (1972) , J. Organomet. Chem. , 46,255 .
- [5] Fischer A., Morgan M. and Eaborn C., (1977) , J. Organomet. Chem. , 136,323 .
- [6] Eaborn C. and Seconi G., (1976) , J. Chem. Soc. Perkin II 925 .
- [7] Mecciantelli D., Seconi G. and Eaborn C., (1978) , J. Chem. Soc. Perkin II, 834 .
- [8] Eaborn C. and Seconi G., (1979) , J. Chem. Soc. Perkin II, 203 .
- [9] Eaborn C., Stamper J. and Seconi G., (1980) , J. Organomet. Chem 202 , 1878 .
- [10] Eaborn C. and Mahmoud F. (1981) , J. Organomet. Chem. 206 49 .
- [11] Alexander R., Asomaning W., Eaborn C., Jenkins J. and Walton D., (1974) J. Chem. Soc. Perkin II , 490 .
- [12] Eaborn C., Walton D. and Seconi G., (1976) , J. Chem. Soc. Perkin II , 1857 .
- [13] Eaborn C., Walton D. and Seconi G., (1975) , J. Chem. Soc. Chem. Com., 937 .
- [14] Bender M. and Komiyama , (1978) , Cyclodextrin Chemistry , Springer-Verlag PP. 9 .
- [15] Bender M. and komiyama , (1978) , ibid. PP . 33 .
- [16] Bender M. and Komiyama M., (1977) , J. Am. Chem. Soc. 99, 8021 .
- [17] Van Elten R., Clowes G., Sebastian J. and Bender M., (1967) , J. Am. Chem. Soc., 89 , 3253 .
- [18] Chin T., Chung P. and Lach J., (1968) , J. Pharm. Sci., 57, 44.
- [19] Van Hooijdonk C. and Groos C., (1970) , Rec. Trav. Chim. , 89, 845 .
- [20] Thorna J. and Stewart L., (1965) , Starch , Chemistry and Technology , Whistler R., Paschall (eds .) N.Y. Acad. Press . PP . 209–249 .
- [21] Van Etten R., Sebastian J., Clowes G. and Bender M., (1967) , J. Am. Chem. Soc. , 89 , 32–42 .
- [22] Reference 14 PP. 35–36 .
- [23] Eaborn C. and Parker S., (1954) , J. Chem. Soc. 939 .
- [24] Bott R., Eaborn C., Pande K. and Swaddle T., (1962) , J. Chem. Soc., 1217 .